

THE COMPLEX-SCALED HALF-SPACE MATCHING METHOD*

ANNE-SOPHIE BONNET-BEN DHIA[†], SIMON N. CHANDLER-WILDE[‡], SONIA FLISS[†],
CHRISTOPHE HAZARD[†], KARL-MIKAEL PERFEKT[§], AND
YOHANES TJANDRAWIDJAJA[¶]

Abstract. The half-space matching (HSM) method has recently been developed as a new method for the solution of two-dimensional scattering problems with complex backgrounds, providing an alternative to perfectly matched layers or other artificial boundary conditions. Based on half-plane representations for the solution, the scattering problem is rewritten as a system of integral equations in which the unknowns are restrictions of the solution to the boundaries of a finite number of overlapping half-planes contained in the domain: this integral equation system is coupled to a standard finite element discretization localized around the scatterer. While satisfactory numerical results have been obtained for real wavenumbers, well-posedness and equivalence to the original scattering problem have been established only for complex wavenumbers. In the present paper, by combining the HSM framework with a complex-scaling technique, we provide a new formulation for real wavenumbers which is provably well-posed and has the attraction for computation that the complex-scaled solutions of the integral equation system decay exponentially at infinity. The analysis requires the study of double-layer potential integral operators on intersecting infinite lines, and their analytic continuations. The effectiveness of the method is validated by preliminary numerical results.

Key words. Helmholtz equation, scattering, integral equation, artificial radiation condition

AMS subject classifications. 35J05, 35J25, 35P25, 45B05, 45F15, 65N30, 65N38, 78A45

DOI. 10.1137/20M1387122

1. Introduction and the scattering problem. The mathematical and numerical analysis of scattering by bounded obstacles and/or inhomogeneities in a homogeneous background is a mature research area, and there are many effective numerical schemes, at least for low to moderately high frequencies. However, when the background is heterogeneous (stratified, periodic, etc.) and/or anisotropic, especially when electromagnetic or elastic waves are considered, many theoretical questions are still open and the design of efficient numerical methods remains a significant challenge.

It is well known that, for homogeneous backgrounds, the Sommerfeld radiation condition ensures well-posedness of the scattering problem [25]. The extension of this standard radiation condition to the aforementioned more complex backgrounds is really intricate (see, e.g., [10, 13, 29, 33, 36]). Moreover, a Green's function or tensor, which could be used to derive an integral equation formulation of the problem, is in

*Received by the editors December 21, 2020; accepted for publication (in revised form) September 7, 2021; published electronically January 18, 2022.

<https://doi.org/10.1137/20M1387122>

Funding: The work of the fifth author was partially supported by UK Engineering and Physical Sciences Research Council (EPSRC) grant EP/S029486/1. The work of the sixth author was supported by the French Commissariat à l'Energie Atomique et aux Energies Alternatives (CEA), through funding for Ph.D. studies.

[†]POEMS (CNRS-INRIA-ENSTA Paris), Institut Polytechnique de Paris, 91764 Palaiseau, France (anne-sophie.bonnet-bendhia@ensta-paris.fr, sonia.fliss@ensta-paris.fr, christophe.hazard@ensta-paris.fr).

[‡]Department of Mathematics and Statistics, University of Reading, Whiteknights PO Box 220, Reading RG6 6AX, UK (s.n.chandler-wilde@reading.ac.uk).

[§]Department of Mathematical Sciences, Norwegian University of Science and Technology (NTNU), NO-7491 Trondheim, Norway (karl-mikael.perfekt@ntnu.no).

[¶]Lehrstuhl I, Fakultät für Mathematik, Technische Universität Dortmund, Vogelpothsweg 87, 44227 Dortmund, Germany (Yohanes.Tjandrawidjaja@math.tu-dortmund.de).

general not available or hard to compute. Finally, perfectly matched layer (PML) techniques, which are popular in homogeneous backgrounds because they are easy to implement, can produce spurious effects for complex backgrounds, as is well known for anisotropic backgrounds in relation to instabilities in the time domain [8].

To overcome these difficulties a new method, called the half-space matching (HSM) method, has been developed as an (exact) artificial boundary condition for two-dimensional (2D) scattering problems. This method is based on explicit or semiexplicit expressions for the outgoing solutions of radiation problems in half-planes, these expressions established by using Fourier, generalized Fourier, or Floquet transforms when the background is, respectively, homogeneous [12, 11] (and possibly anisotropic [46, 7, 45]), stratified [42], or periodic [32]. The domain exterior to a bounded region enclosing the scatterers is covered by a finite number of half-planes (at least three). The unknowns of the formulation are the traces of the solution on the boundaries of these half-planes and the restriction of the solution to the bounded region. The system of equations which couples these unknowns is derived by writing compatibility conditions between the different representations of the solution. This coupled system includes second-kind integral equations on the infinite boundaries of the half-planes.

This new formulation is attractive and versatile as a method to truncate computational domains in problems of scattering by localized inhomogeneities in complex backgrounds (including backgrounds that may be different at infinity in different directions). It has been employed successfully in numerical implementations for a range of problems, namely periodic [34] and stratified media (including cases with different stratifications in different parts of the background domain) [42], and anisotropic scalar and elastic problems [46, 45].

Up to now the theoretical and numerical analysis of the method has remained an open question in the challenging, and more practically relevant, nondissipative case when waves radiate out to infinity. But a rather complete analysis has been carried out in the simpler dissipative case, when the solution (and its traces) decay exponentially at infinity. In that case the analysis can be done using an L^2 framework for the traces, and the associated formulation has been shown to be of Fredholm type and well-posed in a number of cases [12, 11], with the sesquilinear form of the weak formulation coercive plus compact, enabling standard numerical analysis arguments [11]. One of the main difficulties in the nondissipative case is the slow decay at infinity of the solution which results in non- L^2 traces. The possibility, to address this, of working in the framework initially introduced in [14, 15] was investigated by the authors, but it seems to be inappropriate for the present analysis.

The objective of this paper is to propose a new formulation of the HSM method which is well-suited for theoretical and numerical analysis (and practical computation) in the nondissipative case. For the sake of clarity and as a first step, we restrict ourselves in the present paper to a relatively simple model problem for which the justification of the method is based on the simple form of the associated Green's function. (Let us mention that the extension of this formulation to anisotropic backgrounds has been already validated and will be the subject of a forthcoming paper. See section 8 for a brief discussion of extensions to other more complex configurations.) This new formulation exploits a fundamental property of the solution in the spirit of the ideas behind complex-scaling methods (e.g., the pioneering works of [6], [2]): in any given direction, the solution, as a function of the associated curvilinear abscissa, has an analytic continuation into the complex plane which is exponentially decaying in the upper complex plane. This enables us to replace the system of equations for the traces by similar equations for exponentially decaying analytical continuations of these

traces. This recovers well-posedness in an L^2 framework, with coercive plus compact sesquilinear forms; moreover, attractive for computation, the rate of exponential decay of these analytically continued traces increases with increasing wavenumber. Let us mention that in [39] a similar integral-equation-based complex-scaling idea, namely a boundary integral equation formulation of PML, is used to compute 2D scattering by localized perturbations in a straight interface between different media.

In the present paper we consider the rather simple model case of a scalar equation, the isotropic Helmholtz equation

$$(1.1) \quad -\Delta u - k^2 \rho u = f \text{ in } \Omega,$$

deduced from the wave equation assuming a time-dependence $e^{-i\omega t}$ for a given angular frequency $\omega > 0$. Here ρ is a function in $L^\infty(\Omega)$, bounded from below by a positive constant, and such that $\rho - 1$ is compactly supported, and the constant $k = \omega/c$ is the wavenumber, where c is the wave speed outside the support of $\rho - 1$, so that (1.1) models propagation in a domain with a local perturbation in wave speed.

The propagation domain Ω is \mathbb{R}^2 , or \mathbb{R}^2 minus a set of obstacles which are included in a bounded region. We assume that, for a positive constant a ,

$$\partial\Omega \subset \bar{\Omega}_a \text{ where } \Omega_a := (-a, a)^2.$$

In the presence of obstacles, boundary conditions have to be added to the model. The source term f is supposed to be a function in $L^2(\Omega)$ with compact support included in Ω_a , and we assume that

$$\text{supp}(\rho - 1) \subset \Omega_a.$$

As already mentioned, in order to get well-posedness, one has to prescribe in addition the Sommerfeld radiation condition, that, for $\mathbf{x} := (x_1, x_2) \in \mathbb{R}^2$,

$$(1.2) \quad \frac{\partial u}{\partial r}(\mathbf{x}) - iku(\mathbf{x}) = o\left(r^{-1/2}\right) \quad \text{as } r := |\mathbf{x}| \rightarrow +\infty,$$

uniformly with respect to $\hat{\mathbf{x}} := \mathbf{x}/r$.

In what follows, we will consider two configurations in order to focus first on the HSM formulation and then on its coupling with a variational formulation in a bounded region. In sections 2 to 5 we consider a Dirichlet problem set in the exterior of the square Ω_a . The application of the analysis in sections 2–5 to general configurations, with source terms, inhomogeneities, and/or obstacles contained in Ω_a , is the object of section 6.

The outline of the paper is as follows. In section 2 we recall the main results concerning the HSM formulation in the dissipative case (that is, with a complex wavenumber k). In previous papers the HSM formulation has been derived using Fourier representations for the solution in half-planes. Here we introduce a new formulation using double-layer potential integral representations to derive the so-called complex-scaled version of the method, valid for a real wavenumber k .

The derivation and the analysis of this new formulation are the object of section 3, which finishes with a statement of our main well-posedness result for the new method. In section 4 we establish properties of the solution that can be reconstructed a posteriori, from knowledge of the complex-scaled traces, notably elements of the far-field pattern. These properties are used in section 5 to establish the uniqueness result for the complex-scaled HSM problem.

The HSM method approach is extended to general configurations in section 6; the analysis in this section depends throughout on the well-posedness and other results

obtained in sections 3–5. In section 7 the implementation of a finite element discretization of the complex-scaled HSM formulation is described and numerical results are presented, for both the model Dirichlet problem and more general configurations. The paper finishes with three appendices to which we defer certain technical details of the analysis.

2. The HSM method for complex wavenumber. In this section, as preparation for studying the HSM method for real wavenumber, we first recall what is known about the method in the dissipative case. It is enough for this purpose to consider the Dirichlet problem for complex wavenumber ($\Im(k) > 0$, $\Re(k) > 0$) in the exterior of the square Ω_a , i.e.,

$$(2.1) \quad \begin{cases} -\Delta u - k^2 u = 0 & \text{in } \Omega := \mathbb{R}^2 \setminus \overline{\Omega_a}, \\ u = g & \text{on } \Sigma_a := \partial\Omega_a, \end{cases}$$

for a given $g \in H^{1/2}(\Sigma_a)$. It is well-known that problem (2.1) has a unique solution $u \in H^1(\Omega)$.

Let us first recall the main results of [12].

The domain Ω is the union of four overlapping half-planes Ω^j that abut the four edges of the square Ω_a . We introduce the following local coordinates for all $j \in \llbracket 0, 3 \rrbracket := \{0, 1, 2, 3\}$:

$$(2.2) \quad \begin{pmatrix} x_1^j \\ x_2^j \end{pmatrix} := \begin{pmatrix} \cos(j\pi/2) & \sin(j\pi/2) \\ -\sin(j\pi/2) & \cos(j\pi/2) \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}.$$

The half-planes and their boundaries are defined as follows for all $j \in \llbracket 0, 3 \rrbracket$:

$$(2.3) \quad \Omega^j := \{(x_1^j, x_2^j) : x_1^j > a, x_2^j \in \mathbb{R}\}, \quad \Sigma^j := \{(x_1^j, x_2^j) : x_1^j = a, x_2^j \in \mathbb{R}\}.$$

Finally, we denote

$$(2.4) \quad \Sigma_a^j := \Sigma_a \cap \Sigma^j.$$

These notations are summarized in Figure 2.1. As explained in the introduction, the formulation uses the representation of the solution in each half-plane Ω^j in terms of its trace on Σ^j . More precisely, let us denote

$$(2.5) \quad \varphi^j := u|_{\Sigma^j} \quad \text{for } j \in \llbracket 0, 3 \rrbracket$$

so that

$$(2.6) \quad u|_{\Omega^j} = U^j(\varphi^j) \quad \text{for } j \in \llbracket 0, 3 \rrbracket,$$

where, for any $\psi \in H^{1/2}(\Sigma^j)$, $U^j(\psi) \in H^1(\Omega^j)$ is the unique solution of

$$(2.7) \quad \begin{cases} -\Delta U^j - k^2 U^j = 0 & \text{in } \Omega^j, \\ U^j = \psi & \text{on } \Sigma^j. \end{cases}$$

In what follows, we identify any function defined on Σ^j , in particular the function φ^j , with a function of the real variable x_2^j .

We can express $U^j(\psi)$ explicitly in terms of its trace ψ in two manners: using the Fourier transform or using a Green's function representation. First, using the Fourier




FIG. 2.1. The notations defined in (2.2)–(2.3)–(2.4).

transform in the x_2^j -direction, it is easy to see that the solution of (2.7) is given by

$$(2.8) \quad U^j(\psi)(\mathbf{x}^j) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \widehat{\psi}(\xi) e^{-\sqrt{\xi^2 - k^2}(x_1^j - a)} e^{i\xi x_2^j} d\xi, \quad \mathbf{x}^j := (x_1^j, x_2^j) \in \Omega^j,$$

where the square root is defined with the convention $\Re(\sqrt{z}) \geq 0$, for $z \in \mathbb{C} \setminus \mathbb{R}^-$ (with $\mathbb{R}^- := (-\infty, 0]$) and $\widehat{\psi}$ is the Fourier transformation of ψ using the convention

$$(2.9) \quad \widehat{\psi}(\xi) := \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \psi(x_2^j) e^{-i\xi x_2^j} dx_2^j, \quad \xi \in \mathbb{R}.$$

Second, using a Green's function representation, we can show that

$$(2.10) \quad U^j(\psi)(\mathbf{x}^j) = \int_{\Sigma^j} \frac{\partial G^j(\mathbf{x}^j, \mathbf{y}^j)}{\partial n(\mathbf{y}^j)} \psi(\mathbf{y}^j) ds(\mathbf{y}^j), \quad \mathbf{x}^j \in \Omega^j,$$

where $G^j(\mathbf{x}^j, \mathbf{y}^j)$ is the Dirichlet Green's function for Ω^j and $n(\mathbf{y}^j)$ is the unit normal to Σ^j that points into Ω^j . Explicitly, $G^j(\mathbf{x}^j, \mathbf{y}^j) = \Phi(\mathbf{x}^j, \mathbf{y}^j) - \Phi(\tilde{\mathbf{x}}^j, \mathbf{y}^j)$, with $\tilde{\mathbf{x}}^j$ the image of \mathbf{x}^j in Σ^j , where $\Phi(\mathbf{x}, \mathbf{y})$ is the standard fundamental solution of the Helmholtz equation defined by

$$(2.11) \quad \Phi(\mathbf{x}, \mathbf{y}) := \frac{i}{4} H_0^{(1)}(k|\mathbf{x} - \mathbf{y}|), \quad \mathbf{x}, \mathbf{y} \in \mathbb{R}^2, \quad \mathbf{x} \neq \mathbf{y},$$

so that, equivalently,

$$(2.12) \quad U^j(\psi)(\mathbf{x}^j) = 2 \int_{\Sigma^j} \frac{\partial \Phi(\mathbf{x}^j, \mathbf{y}^j)}{\partial n(\mathbf{y}^j)} \psi(\mathbf{y}^j) ds(\mathbf{y}^j), \quad \mathbf{x}^j \in \Omega^j.$$

This leads to

$$(2.13) \quad U^j(\psi)(\mathbf{x}^j) = \int_{\mathbb{R}} h(x_1^j - a, x_2^j - y_2^j) \psi(y_2^j) dy_2^j, \quad \mathbf{x}^j \in \Omega^j,$$




FIG. 2.2. Construction of the compatibility condition.

where

$$(2.14) \quad h(x_1, x_2) := \frac{ikx_1}{2} \frac{H_1^{(1)}(kR(x_1, x_2))}{R(x_1, x_2)},$$

and

$$(2.15) \quad R(x_1, x_2) := (x_1^2 + x_2^2)^{1/2}, \quad x_1, x_2 \in \mathbb{R}.$$

Let us remark that the two representations (2.8) and (2.13) of $U^j(\psi)$ can be derived one from other by using simply a Plancherel formula (e.g., [19, p. 821]).

To derive the system of equations whose unknowns are the traces φ^j of the solution, it suffices to write that the half-plane representations must coincide where they coexist. For instance, in the quarter plane $\Omega^0 \cap \Omega^1$ (see Figure 2.2) we have

$$(2.16) \quad u = U^0(\varphi^0) = U^1(\varphi^1) \quad \text{in } \Omega^0 \cap \Omega^1,$$

and in particular

$$(2.17) \quad \varphi^1 = U^0(\varphi^0) \quad \text{on } \Omega^0 \cap \Sigma^1,$$

which leads to a first integral equation linking φ^0 and φ^1 . Indeed, for any point of $\Sigma^1 \cap \Omega^0$, represented by (x_1^0, x_2^0) and (x_1^1, x_2^1) in local coordinates systems, we have $x_2^1 = -x_1^0 < -a$ and $x_1^1 = x_2^0 = a$ (indeed, the point is on $\Sigma^1 \cap \Omega^0$ if and only if these equations hold). Thus the compatibility relation (2.17) can be rewritten (identifying Σ^1 with \mathbb{R} in the way noted above) as

$$(2.18) \quad \varphi^1(x_2^1) = U^0(\varphi^0)(-x_2^1, a), \quad x_2^1 < -a,$$

where we can use either of the two integral representations (2.8) and (2.13) for the half-plane solution $U^0(\varphi^0)$. From (2.16) we have also

$$\varphi^0 = U^1(\varphi^1) \quad \text{on } \Sigma^0 \cap \Omega^1,$$

which leads to another integral equation linking φ^0 and φ^1 :

$$\varphi^0(x_2^0) = U^1(\varphi^1)(x_2^0, -a), \quad x_2^0 > a.$$

Repeating this for each quarter plane we get eight equations linking the four traces. In order to write the system of equations in a condensed manner, we shall use the same notation t instead of the various variables x_2^j so that the eight equations become

$$(2.19) \quad \forall j \in \llbracket 0, 3 \rrbracket, \quad \begin{cases} \varphi^j(t) = S D \varphi^{j-1}(t), & t < -a, \\ \varphi^j(t) = D S \varphi^{j+1}(t), & t > a, \end{cases}$$

where we have set $\varphi^{-1} := \varphi^3$ and $\varphi^4 := \varphi^0$, and where the operators S and D are defined as follows. For any $\psi \in L^2(\mathbb{R})$,

$$(2.20) \quad S\psi(t) := \psi(-t), \quad t \in \mathbb{R},$$

and the integral operator D is defined by

$$(2.21) \quad D\psi(t) := U^0(\psi)(t, a), \quad t > a.$$

D can be given explicitly by either of the following two expressions:

$$(2.22) \quad D\psi(t) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \widehat{\psi}(\xi) e^{i\sqrt{k^2 - \xi^2}(t-a)} e^{i\xi a} d\xi, \quad t > a,$$

or

$$(2.23) \quad D\psi(t) = \int_{\mathbb{R}} h(t-a, s-a) \psi(s) ds, \quad t > a,$$

where the kernel h is defined in (2.14).¹ The system of equations has to be completed with the Dirichlet boundary condition rewritten as

$$(2.24) \quad \varphi^j(t) = g|_{\Sigma_a^j}(t), \quad -a < t < a, \quad j \in \llbracket 0, 3 \rrbracket.$$

One can easily check that (2.19)–(2.24) is equivalent to the original problem (2.1). More precisely, if $\{\varphi^0, \varphi^1, \varphi^2, \varphi^3\} \in (H^{1/2}(\mathbb{R}))^4$ is a solution to (2.19)–(2.24), then one can recover the solution u to (2.1), from the knowledge of the φ^j 's, thanks to the half-plane representations (2.8) or (2.13). Indeed, by uniqueness of Dirichlet quarter-plane problems, two half-plane representations $U^j(\varphi^j)$ and $U^{j+1}(\varphi^{j+1})$ coincide on the quarter-plane $\Omega^j \cap \Omega^{j+1}$ since the compatibility conditions (2.19) imply that they coincide on its boundary.

For the analysis and the computation it is convenient to consider the formulation in an L^2 -framework:

$$(2.25) \quad \begin{array}{l} \text{Find } \{\varphi^0, \varphi^1, \varphi^2, \varphi^3\} \in (L^2(\mathbb{R}))^4 \text{ such that} \\ \left| \begin{array}{ll} \varphi^j(t) = S D \varphi^{j-1}(t), & t < -a, \\ \varphi^j(t) = g|_{\Sigma_a^j}(t), & -a < t < a, \quad j \in \llbracket 0, 3 \rrbracket, \\ \varphi^j(t) = D S \varphi^{j+1}(t), & t > a. \end{array} \right. \end{array}$$

One attraction of this L^2 -framework is that it allows the use of elementary operations of restriction and extension. More precisely, for any function of $L^2(\mathbb{R})$, its restriction to an open interval $I \subset \mathbb{R}$ is in $L^2(I)$. More significantly, any function of $L^2(I)$

¹Note that, as is clear from (2.12), D is precisely a double-layer potential operator (in the sense, e.g., of [26] or [17]) from $\{(a, t) : t \in \mathbb{R}\}$ to $\{(t, a) : t \geq a\}$.

extended by 0 belongs to $L^2(\mathbb{R})$. For simplicity, any function defined on a part of \mathbb{R} is identified hereafter with its extension by 0. With this convention we can write

$$(2.26) \quad L^2(\mathbb{R}) = L^2(-\infty, -a) \oplus L^2(-a, a) \oplus L^2(a, +\infty),$$

which will be extensively used hereafter. In line with this convention, we define $D\psi(t)$ for all $t \in \mathbb{R}$ by setting

$$(2.27) \quad D\psi(t) := 0, \quad t \leq a.$$

With these various conventions (2.25) can be rewritten briefly as

$$(2.28) \quad \varphi^j = SD\varphi^{j-1} + DS\varphi^{j+1} + g|_{\Sigma_a^j}, \quad j \in \llbracket 0, 3 \rrbracket.$$

Also, with the above conventions, results proved in [17, 12] can be stated as follows.

PROPOSITION 2.1.

- (i) D is a continuous operator on $L^2(\mathbb{R})$, with range in $L^2(a, +\infty) \subset L^2(\mathbb{R})$.
- (ii) As an operator on $L^2(a, +\infty)$, D is the sum of an operator of norm $\leq 1/\sqrt{2}$ and a compact operator.
- (iii) As an operator from $L^2(-\infty, -a)$ to $L^2(a, +\infty)$, D is compact.

Proof. Note first that (i) follows immediately from (ii) and (2.27), since (ii), together with a symmetry argument with respect to a , implies that $D: L^2(-\infty, a) \rightarrow L^2(a, +\infty)$ is also the sum of an operator of norm $\leq 1/\sqrt{2}$ and a compact operator. Consider the expression (2.23) for D . Because of the dissipation ($\Im(k) > 0$), the kernel h is exponentially decaying at infinity (i.e., as t or s tends to ∞). Further, the mapping $(t, s) \mapsto h(t - a, s - a)$ is continuous except at $t = s = a$. Thus (iii) is clear since the kernel of D is Hilbert–Schmidt, i.e.,

$$(t, s) \mapsto h(t - a, s - a) \in L^2((-\infty, -a) \times (a, +\infty)).$$

To show (ii), the only difficulty comes from the singularity of the kernel h at $t = s = a$. As in Proposition C.2 and Remark C.3 in Appendix C, let h_0 and D_0 denote h and D , respectively, when $k = 0$. For $b > a$ let $\chi_{(a,b)}$ denote the characteristic function of (a, b) . Then it is straightforward to see that, for every $b > a$, $D - \chi_{(a,b)}D_0$ is an integral operator with kernel $h(t - a, s - a) - \chi_{(a,b)}(t)h_0(t - a, s - a)$ that is Hilbert–Schmidt, so $D - \chi_{(a,b)}D_0$ is compact. Further (Remark C.3), as an operator on $L^2(a, +\infty)$, $\|D_0\| = 1/\sqrt{2}$, so also $\|\chi_{(a,b)}D_0\| \leq \|\chi_{(a,b)}\| \|D_0\| = 1/\sqrt{2}$. \square

Now the system (2.25) can be formulated in an operator form. Let us introduce

$$(2.29) \quad L_0^2(\mathbb{R}) := \{\psi \in L^2(\mathbb{R}) : \psi(t) = 0 \text{ for } -a < t < a\}$$

and

$$\Phi := \{\varphi^0, \varphi^1, \varphi^2, \varphi^3\} \in (L^2(\mathbb{R}))^4,$$

$$(2.30) \quad \Phi_g := \{g|_{\Sigma_a^0}, g|_{\Sigma_a^1}, g|_{\Sigma_a^2}, g|_{\Sigma_a^3}\} \in (L^2(-a, a))^4 \subset (L^2(\mathbb{R}))^4.$$

Then, noting (2.28), the system (2.25) can be rewritten as

$$(2.31) \quad \begin{aligned} &\text{Find } \Phi \in (L^2(\mathbb{R}))^4 \text{ such that } \Phi - \Phi_g \in (L_0^2(\mathbb{R}))^4 \text{ and} \\ &(\mathbb{I} - \mathbb{D})(\Phi - \Phi_g) = \mathbb{D}\Phi_g, \end{aligned}$$

where

$$(2.32) \quad \mathbb{D} := \begin{bmatrix} 0 & DS & 0 & SD \\ SD & 0 & DS & 0 \\ 0 & SD & 0 & DS \\ DS & 0 & SD & 0 \end{bmatrix}.$$

In [12], the following result is proven.

THEOREM 2.2.

- (i) \mathbb{D} is a continuous operator on $(L^2(\mathbb{R}))^4$ and $\mathbb{D}((L^2(\mathbb{R}))^4) \subset (L_0^2(\mathbb{R}))^4$.
- (ii) As an operator on $(L_0^2(\mathbb{R}))^4$, \mathbb{D} is the sum of an operator of norm $\leq 1/\sqrt{2}$ and a compact operator.
- (iii) Problem (2.31) is well-posed.

Let us give some ideas of the proof which will be relevant for the following sections. The property (iii) is largely a consequence of (ii) since (ii) gives that $(\mathbb{I} - \mathbb{D})$, as an operator acting on $(L_0^2(\mathbb{R}))^4$, is the sum of a coercive operator² and a compact one. By Fredholm theory, it suffices then to show uniqueness (which is not straightforward in the L^2 framework; see [12] and also [9] for more details). The properties (i) and (ii) are consequences (see Appendix A) of Proposition 2.1.

3. The complex-scaled HSM method for real wavenumber. In this section we consider the Dirichlet problem of the previous section, but now with real wavenumber ($k > 0$). Where $H_{\text{loc}}^1(\Omega) := \{v|_{\Omega} : v \in H_{\text{loc}}^1(\mathbb{R}^2)\}$, we seek $u \in H_{\text{loc}}^1(\Omega)$ such that

$$(3.1) \quad \begin{cases} -\Delta u - k^2 u = 0 & \text{in } \Omega := \mathbb{R}^2 \setminus \overline{\Omega_a}, \\ u = g & \text{on } \Sigma_a, \end{cases}$$

for a given $g \in H^{1/2}(\Sigma_a)$, and such that the radiation condition (1.2) holds. It is well known that this problem has a unique solution.

The half-plane representations (2.6), with $U^j(\varphi^j)$ given by (2.10) (equivalently (2.12) or (2.13)), still hold for k real and can be derived via Green's theorem using the radiation condition (1.2) (cf. [18, Theorem 2.1]). As a consequence, the traces φ_j , $j \in \llbracket 0, 3 \rrbracket$, of the solution u on Σ^j , still satisfy the system of equations (2.25) when $k > 0$. We note that, although the solution of (3.1)–(1.2) decays only slowly, like $r^{-1/2}$ as $r \rightarrow +\infty$, the integrals (2.13) still make sense. The bound (C.1), which follows from asymptotics of the Hankel function $H_1^{(1)}$, implies that, for some constant $C > 0$ depending only on k ,

$$|h(x_1, x_2)| \leq Cx_1(R^{-3/2} + R^{-2}), \quad x_1 > 0, \quad x_2 \in \mathbb{R},$$

where $R := (x_1^2 + x_2^2)^{1/2}$, so that (2.13) is well-defined, even when k is real, for every $\mathbf{x}^j \in \Omega^j$ and $\psi \in L_{\text{loc}}^2(\mathbb{R})$ with $\psi(t) = \mathcal{O}(1)$ as $|t| \rightarrow +\infty$. Moreover, though we shall not need this, it is still possible to rewrite (2.13) equivalently as (2.8), provided that care is taken in interpreting the right-hand side of (2.8); see the discussion in [4, 20] and [14, 15].

From a numerical point of view, the HSM method for real k works well [12]. However, from a theoretical point of view, the formulation does not make sense in

²Recall that, given a Hilbert space \mathcal{H} with inner product (\cdot, \cdot) , we call a bounded linear operator A on \mathcal{H} *coercive* if the corresponding sesquilinear form $a(\cdot, \cdot)$, defined by $a(\phi, \psi) = (A\phi, \psi)$ for all $\phi, \psi \in \mathcal{H}$, is coercive, i.e., if, for some constant $\gamma > 0$, $\Re(a(\phi, \phi)) \geq \gamma\|\phi\|^2$ for all $\phi \in \mathcal{H}$.

an L^2 setting. Indeed, as the solution of (3.1)–(1.2) decays only like $r^{-1/2}$ as $r \rightarrow +\infty$, we cannot expect that its traces on Σ^j , $j \in \llbracket 0, 3 \rrbracket$, are in $L^2(\mathbb{R})$. In parallel work [9] we have shown that the HSM formulation for real k is equivalent with the original problem (3.1) if we supplement it with radiation conditions analogous to the Sommerfeld condition (1.2) (see (5.1) below). However, there are still significant gaps in our understanding of this formulation when k is real. In particular, while (2.25) can be written formally in operator form as (2.31), just as in the dissipative case, in the case when k is real there is no obvious function space setting (e.g., L^p , or a weighted L^p space) for which this formulation makes sense, with \mathbb{D} a well-defined bounded linear operator. Consequently, we are not able to justify the numerical method and neither provide a priori error estimates.

These difficulties with the standard formulation for real k are part of the motivation for the method proposed in this paper that we term the *complex-scaled HSM method*. The idea behind this method is to “complexify.” Since the pioneering works of Aguilar, Balslev, and Combes [2, 6], complex-scaling methods have been used intensively to construct the analytic continuation of resolvents in mathematical physics; see, for instance, [31] and the references therein. The complex-scaling method in [2, 6] is closely related to the idea behind PML (see, for instance, [24]). Our use of complex-scaling is somewhat different since we consider only analytic continuation of traces of the solution on particular infinite half-lines and apply that complex scaling in an integral equation context. This is similar to manipulations made to understand analyticity of boundary traces in high frequency scattering problems in [22, section 4.1]. Precisely our plan is as follows:

1. From properties of the solution u of (3.1)–(1.2) we deduce that the traces φ^j , $j \in \llbracket 0, 3 \rrbracket$, have analytic continuations into the complex plane from $(-\infty, -a)$ and from $(a, +\infty)$. Further, we introduce paths in the complex plane on which the φ^j ’s are L^2 (in fact, decay exponentially); see Proposition 3.3. The objective of the next steps is to derive an equivalent of the HSM formulation for these “complex-scaled” traces.
2. For real wavenumbers (2.6) and (2.13) provide half-plane representations of the solution u in terms of the traces φ^j , $j \in \llbracket 0, 3 \rrbracket$. The magic result is that the solution u can also be represented in terms of the complex-scaled traces; see Theorem 3.4. The price to pay is that the j th representation, for $j \in \llbracket 0, 3 \rrbracket$, is valid only in a part, which depends on the path chosen in step 1, of the corresponding half-plane Ω^j . These new representations are deduced from the initial ones (2.6) by applying Cauchy’s integral theorem.
3. Fortunately, the part of Ω^j where the new representation holds contains the half-lines $\Sigma^{j\pm 1} \cap \Omega^j$. By complexifying—by which we mean analytically continuing—this new representation into the complex plane from the half-lines $\Sigma^{j\pm 1} \cap \Omega^j$, we can derive compatibility conditions for the complex-scaled traces which constitute a complex-scaled version of the HSM formulation in an L^2 setting. Fredholmness of this formulation can be proven using similar arguments as for the standard HSM for complex wavenumbers; see Theorem 3.7.
4. Once the complex-scaled traces are computed the solution can be reconstructed using the new representations in terms of the complex-scaled traces established in step 2.

Let us mention that, while the initial motivation for the complexification was a theoretical one, it turns out that the new formulation is very attractive computationally, because of the fast decay at infinity of the complex-scaled traces. Let us

note also that this idea of complexification is potentially valuable for computation in the dissipative case too, and it is likely that the formulation in the nondissipative case could be derived from a complex-scaled formulation in the dissipative case by a limiting absorption argument. (Something similar has been done in the context of scattering by wedges in [28, 37].) An attraction of such a derivation would be that traces are in L^2 at each step of the derivation.

3.1. The complex-scaled traces. The construction of the so-called complex-scaled traces is based on an analyticity property of any solution $u \in H_{\text{loc}}^1(\Omega)$ to (1.2)–(3.1), which can be derived as follows. Following [16] (and see [21, Theorem 2.27, Corollary 2.28]), u can be expressed as a combined single- and double-layer potential on Σ_a , i.e., as

$$(3.2) \quad u(\mathbf{x}) = \mathcal{D}\phi(\mathbf{x}) - ik\mathcal{S}\phi(\mathbf{x}), \quad \mathbf{x} \in \Omega,$$

for some $\phi \in H^{1/2}(\Sigma_a)$ (specified below). Here $\mathcal{S}\phi$ and $\mathcal{D}\phi$ are the (acoustic) single- and double-layer potentials, respectively, with density ϕ , defined for $\phi \in L^2(\Sigma_a)$ by

$$\mathcal{S}\phi(\mathbf{x}) := \int_{\Sigma_a} \Phi(\mathbf{x}, \mathbf{y}) \phi(\mathbf{y}) \, ds(\mathbf{y}), \quad \mathcal{D}\phi(\mathbf{x}) := \int_{\Sigma_a} \frac{\partial \Phi(\mathbf{x}, \mathbf{y})}{\partial n(\mathbf{y})} \phi(\mathbf{y}) \, ds(\mathbf{y}), \quad \mathbf{x} \in \Omega,$$

where the normal n is directed into Ω and Φ is the outgoing Green's function of the Helmholtz equation given in (2.11). The function u defined in (3.2) satisfies the Helmholtz equation (3.1) and the Sommerfeld radiation condition (1.2) for any choice of $\phi \in H^{1/2}(\Sigma_a)$ (in fact, any $\phi \in L^2(\Sigma_a)$) and (see [21, section 2.6]) satisfies the boundary condition $u = g$ on Σ_a provided

$$(3.3) \quad A\phi = g,$$

where $A\phi$ is defined for $\phi \in L^2(\Sigma_a)$ and almost all $\mathbf{x} \in \Sigma_a$ by

$$A\phi(\mathbf{x}) := \frac{\phi(\mathbf{x})}{2} + \int_{\Sigma_a} \left[\frac{\partial \Phi(\mathbf{x}, \mathbf{y})}{\partial n(\mathbf{y})} - ik\Phi(\mathbf{x}, \mathbf{y}) \right] \phi(\mathbf{y}) \, ds(\mathbf{y}),$$

with the integral understood as a Cauchy principal value. Since $A : H^s(\Sigma_a) \rightarrow H^s(\Sigma_a)$ is invertible for $0 \leq s \leq 1$ [23, Corollary 2.8], in particular for $s = 1/2$, (3.3) has a unique solution $\phi \in H^{1/2}(\Sigma_a)$.

For a given $j \in \llbracket 0, 3 \rrbracket$, we apply (3.2) for $\mathbf{x} \in \Sigma^j \setminus \Sigma_a^j$, and we use the coordinate system (x_1^j, x_2^j) defined in (2.2). Defining $\mathbf{x}^j(t) := (a, t)$ with $|t| > a$, this yields, by definition of φ^j ,

$$(3.4) \quad \varphi^j(t) = \mathcal{D}^j\phi(t) - ik\mathcal{S}^j\phi(t),$$

for real t with $|t| > a$, where

$$(3.5) \quad \mathcal{D}^j\phi(t) := \int_{\Sigma_a} \frac{\partial \Phi(\mathbf{x}^j(t), \mathbf{y}^j)}{\partial n(\mathbf{y}^j)} \phi(\mathbf{y}^j) \, ds(\mathbf{y}^j), \quad \mathcal{S}^j\phi(t) := \int_{\Sigma_a} \Phi(\mathbf{x}^j(t), \mathbf{y}^j) \phi(\mathbf{y}^j) \, ds(\mathbf{y}^j),$$

and we recall from (2.11) that

$$\Phi(\mathbf{x}^j(t), \mathbf{y}^j) = \frac{i}{4} H_0^{(1)}(kR(a - y_1^j, t - y_2^j)),$$

where R is defined in (2.15).

Let us use (3.4) to prove that the function φ^j , defined by (2.5), can be continued analytically into the complex plane from $(-\infty, -a)$ and from $(a, +\infty)$. Consider a fixed $\mathbf{y}^j \in \Sigma_a$. The function $z \mapsto R(a - y_1^j, z - y_2^j)$ has an analytic continuation from $(a, +\infty)$ (respectively, $(-\infty, -a)$) to the complex half-plane $\Re(z) > a$ (respectively, $\Re(z) < -a$). Indeed, to obtain this analytic continuation we simply have to use, in the definition (2.15) of $R(a - y_1^j, z - y_2^j)$ for real z , the principal square root of a complex number, which we will denote by $z^{1/2}$ or \sqrt{z} , defined as

$$\sqrt{z} := |z|^{1/2} e^{i\text{Arg}(z)/2} \quad \text{with } \text{Arg}(z) \in (-\pi, +\pi],$$

which is analytic in $\mathbb{C} \setminus \mathbb{R}^-$, where $\mathbb{R}^- := (-\infty, 0]$. The analyticity of $z \mapsto R(a - y_1^j, z - y_2^j)$ follows by noticing that $(y_1^j - a)^2 + (y_2^j - z)^2 \in \mathbb{C} \setminus \mathbb{R}^-$ if $\Re(z) > a$ (respectively, if $\Re(z) < -a$), since $y_2^j \in [-a, +a]$. Since also $z \mapsto H_0^{(1)}(z)$ is analytic in $\Re(z) > 0$, we conclude that the function $z \mapsto \Phi(\mathbf{x}^j(z), \mathbf{y}^j)$ is analytic in $|\Re(z)| > a$. And the same arguments and conclusion apply also for $z \mapsto \partial\Phi(\mathbf{x}^j(z), \mathbf{y}^j)/\partial n(\mathbf{y}^j)$. Finally, using standard results about analyticity of functions defined as integrals (e.g., [3, Corollary X.3.19]), we conclude that $z \mapsto \mathcal{S}^j\phi(z)$, $z \mapsto \mathcal{D}^j\phi(z)$, and so also $z \mapsto \varphi^j(z)$, have analytic continuations from $(a, +\infty)$ (respectively, $(-\infty, -a)$) to the complex half-plane $\Re(z) > a$ (respectively, $\Re(z) < -a$).

The behavior of these analytic continuations as $|z| \rightarrow +\infty$ depends on $\Im(z)$. Indeed, for $m \in \mathbb{N} := \{0, 1, \dots\}$, we have [1, equation (9.2.30)]

$$(3.6) \quad H_m^{(1)}(z) = \sqrt{\frac{2}{\pi z}} e^{i(z - m\pi/2 - \pi/4)} \left(1 + \mathcal{O}(|z|^{-1})\right) \quad \text{as } |z| \rightarrow +\infty,$$

uniformly in $\text{Arg}(z)$ for $|\text{Arg}(z)| < \zeta$, for every $\zeta < \pi$. Further, as a consequence of Lemma B.2 and Remark B.3,

$$(3.7) \quad R(a - y_1^j, z - y_2^j) = z - y_2^j + \mathcal{O}(|z|^{-1}) \quad \text{in } \Re(z) > a, \quad \text{and}$$

$$(3.8) \quad R(a - y_1^j, z - y_2^j) = y_2^j - z + \mathcal{O}(|z|^{-1}) \quad \text{in } \Re(z) < -a,$$

as $|z| \rightarrow +\infty$, uniformly in $\text{Arg}(z)$ and \mathbf{y}^j , for $\mathbf{y}^j \in \Sigma_a$. Using (3.4) it follows from the above asymptotics that

$$(3.9) \quad \varphi^j(z) = \mathcal{O}\left(e^{-k\Im(z)} |z|^{-1/2}\right) \quad \text{in } \Re(z) > a, \quad \text{and}$$

$$(3.10) \quad \varphi^j(z) = \mathcal{O}\left(e^{k\Im(z)} |z|^{-1/2}\right) \quad \text{in } \Re(z) < -a,$$

as $|z| \rightarrow +\infty$, uniformly with respect to $\text{Arg}(z)$, and we note that $\Im(R(a - y_1^j, z - y_2^j))$ and $\Im(z)$ have the same sign if $\Re(z) > a$ (opposite signs if $\Re(z) < -a$). Thus $\varphi^j(z)$ is exponentially decreasing in the quadrants $\{\Re(z) > a \text{ and } \Im(z) > 0\}$ and $\{\Re(z) < -a \text{ and } \Im(z) < 0\}$.

Our idea is to choose half-lines in these quadrants and consider the analytic continuations of the φ^j on these half-lines as new unknowns instead of the initial traces. In other words, choosing some $\theta \in (0, \pi/2)$, we introduce the complex path (Figure 3.1) parameterized by

$$(3.11) \quad z = \tau_\theta(s) := \begin{cases} -a + (s + a)e^{i\theta} & \text{if } s < -a, \\ s & \text{if } -a \leq s \leq +a, \\ a + (s - a)e^{i\theta} & \text{if } s > a, \end{cases}$$

and we define the complex-scaled traces by

$$(3.12) \quad \varphi_\theta^j(s) := \varphi^j(\tau_\theta(s)) \quad \text{for } s \in \mathbb{R} \text{ and } j \in \llbracket 0, 3 \rrbracket.$$

Remark 3.1. We have chosen particular complex paths in the quadrants $\{\Re(z) > a \text{ and } \Im(z) > 0\}$ and $\{\Re(z) < -a \text{ and } \Im(z) < 0\}$, given by (3.11), that move into the complex plane already from $\pm a$, the corners of Σ_a . Note that it is possible, alternatively, to start to complexify at a positive distance from the corners, i.e., from $\pm b$, for some $b > a$. It is also possible to choose a smoother complex change of variable as usually done in PML methods.




FIG. 3.1. The complex path $s \rightarrow \tau_\theta(s)$.

We note that, by definition, $\varphi_\theta^j|_{(-a,a)} \in L^2(-a,a)$, since $\varphi_\theta^j(t) = \varphi^j(t) = u(\mathbf{x}^j(t)) = g(\mathbf{x}^j(t))$, for $-a < t < a$, and $g \in H^{1/2}(\Sigma_a) \subset L^2(\Sigma_a)$. Moreover, thanks to the analyticity of φ^j , it is clear that the restrictions of φ_θ^j to $(a, +\infty)$ and $(-\infty, -a)$ are continuous. Further, it follows from (3.9)–(3.10) that

$$(3.13) \quad \varphi_\theta^j(s) = \mathcal{O}\left(e^{-k|s|\sin\theta}|s|^{-1/2}\right) \quad \text{as } |s| \rightarrow +\infty.$$

Thus whether or not $\varphi_\theta^j \in L^2(\mathbb{R})$ depends on the behavior of $\varphi_\theta^j(t)$ as $t \rightarrow a^+$ and $t \rightarrow -a^-$.

The following propositions bound $\varphi_\theta^j(t)$ on $|t| > a$, in particular near $\pm a$, and show that $\varphi_\theta^j \in L^2(\mathbb{R})$ for $0 < \theta < \pi/2$. We relegate some of the technical details to Appendix C. Applying the bounds from Proposition C.1 to (the analytic continuation of) (3.4) we obtain the following proposition, on observing, from (3.3), that $\phi = A^{-1}g$ and that [23, Corollary 2.8] A^{-1} is bounded as an operator on $L^2(\Sigma_a)$.

PROPOSITION 3.2. *For every $\theta \in (0, \pi/2)$ and $j \in \llbracket 0, 3 \rrbracket$ there exists a constant $C > 0$, that depends only on a , k , and θ , such that*

$$|\varphi^j(z)| \leq \begin{cases} C|z-a|^{-1/2}e^{-k\Im(z)}\|g\|_{L^2(\Sigma_a)} & \text{if } \Re(z) > a \text{ with } |\arg(z-a)| \leq \theta, \\ C|z+a|^{-1/2}e^{k\Im(z)}\|g\|_{L^2(\Sigma_a)} & \text{if } \Re(z) < -a \text{ with } |\arg(-z-a)| \leq \theta. \end{cases}$$

The bound in the above proposition implies that $\varphi_\theta^j \in L^1(\mathbb{R})$, for $0 < \theta < \pi/2$, and is sufficient for our Cauchy's integral formula arguments below in section 3.2. But it is not quite strong enough to establish $\varphi_\theta^j \in L^2(\mathbb{R})$, as discussed in the proof of Proposition C.4 in Appendix C.

It follows from (the analytic continuation of) (3.4) that, in the notation of Proposition C.4, $\varphi_\theta^j(s) = \mathcal{D}_\theta^j\phi(s) - ik\mathcal{S}_\theta^j\phi(s)$, for $|s| > a$, where $\phi = A^{-1}g$. Thus, and arguing as above Proposition 3.2, we deduce the following result from the above proposition and Proposition C.4.

PROPOSITION 3.3. *For $0 < \theta < \pi/2$ and $j \in \llbracket 0, 3 \rrbracket$, $\varphi_\theta^j \in L^2(\mathbb{R})$. Further, for some constant $C > 0$ depending only on θ , a , and k ,*

$$|\varphi_\theta^j(s)| \leq C(|s|-a)^{-1/2}e^{-k\sin(\theta)(|s|-a)}\|g\|_{L^2(\Sigma_a)}, \quad |s| > a,$$

$$\text{and } \|\varphi_\theta^j\|_{L^2(\mathbb{R})} \leq C\|g\|_{L^2(\Sigma_a)}.$$

3.2. The deformed half-plane representations. We have introduced in the previous section the complex-scaled traces φ_θ^j and proved that they belong to $L^2(\mathbb{R})$. The objective now is to derive an HSM formulation for these new unknowns. The first step is to establish new representation formulas for the solution u of (1.1)–(1.2) in the half-planes using these complex-scaled traces instead of the original traces φ^j .

We recall that the solution u of (1.1)–(1.2) can be represented in each half-plane in terms of its traces as

$$(3.14) \quad u(\mathbf{x}^j) = \int_{\mathbb{R}} h(x_1^j - a, x_2^j - y_2^j) \varphi^j(y_2^j) dy_2^j, \quad \mathbf{x}^j \in \Omega^j,$$

where the kernel h is defined by (2.14). Our objective is to derive a similar formula using the complex-scaled trace φ_θ^j instead of the trace φ^j . This can be done by deforming the path of integration into the complex plane. This leads to the following crucial result.

THEOREM 3.4. *Let u be the solution of (1.1)–(1.2) and φ_θ^j be defined as in (3.12). For $0 < \theta < \pi/2$ we have*

$$(3.15) \quad u(\mathbf{x}^j) = \int_{\mathbb{R}} h(x_1^j - a, x_2^j - \tau_\theta(s)) \varphi_\theta^j(s) \tau_\theta'(s) ds, \quad \mathbf{x}^j \in \Omega_\theta^j, \quad j \in \llbracket 0, 3 \rrbracket,$$

where

$$(3.16) \quad \Omega_\theta^j := \{\mathbf{x}^j = (x_1^j, x_2^j) \in \Omega^j : x_1^j - a > (|x_2^j| - a) \tan(\theta)\}.$$

Remark 3.5. Let us point out that the new representation formula (3.15) is valid only in a subdomain Ω_θ^j of the half-plane Ω^j (Figure 3.2). The larger the angle θ , the faster the decay of the complex-scaled traces φ_θ^j (Proposition 3.3) but the smaller the domain of validity Ω_θ^j of the new representation (3.15).




FIG. 3.2. The domains of validity of the half-plane representations: left is without complex-scaling (Ω^j); right is with complex-scaling (Ω_θ^j).

Proof. To derive (3.15) from (3.14) it suffices to show (note the symmetry $\tau_\theta(-s) = -\tau_\theta(s)$) that, for all $\mathbf{x}^j \in \Omega_\theta^j$,

$$\int_a^{+\infty} h(x_1^j - a, x_2^j - y_2^j) \varphi^j(y_2^j) dy_2^j = \int_a^{+\infty} h(x_1^j - a, x_2^j - \tau_\theta(s)) \varphi_\theta^j(s) \tau_\theta'(s) ds$$

and

$$\int_a^{+\infty} h(x_1^j - a, x_2^j + y_2^j) \varphi^j(-y_2^j) dy_2^j = \int_a^{+\infty} h(x_1^j - a, x_2^j + \tau_\theta(s)) \varphi_\theta^j(-s) \tau_\theta'(s) ds.$$

For $0 < \delta < M$ and $0 < \theta < \pi/2$ we introduce the complex domains

$$T_\theta := \{z = a + re^{i\alpha} : 0 < \alpha < \theta, r > 0\} \quad \text{and} \\ T_\theta^{\delta, M} := \{z = a + re^{i\alpha} : 0 < \alpha < \theta, \delta < r < M\}$$

(see Figure 3.3). First we show that if $\mathbf{x}^j = (x_1^j, x_2^j) \in \Omega^j$, then

$$(3.17) \quad \mathbf{x}^j \in \Omega_\theta^j \Leftrightarrow \begin{cases} z \mapsto h(x_1^j - a, x_2^j \pm z) \text{ is analytic in } T_\theta \\ \text{and continuous in } \overline{T_\theta}. \end{cases}$$

Indeed, for each \mathbf{x}^j , the function $z \mapsto h(x_1^j - a, x_2^j - z)$ has two branch points z_\pm , the points where $(x_1^j - a)^2 + (x_2^j - z)^2$ vanishes, given by $z_\pm = x_2^j \pm i(x_1^j - a)$. If $\mathbf{x}^j = (x_1^j, x_2^j) \in \Omega^j$ these branch points are outside $\overline{T_\theta}$ if and only if $x_1^j - a > (x_2^j - a) \tan(\theta)$; similarly, the branch points of the mapping $z \mapsto h(x_1^j - a, x_2^j + z)$ are outside $\overline{T_\theta}$ if and only if $x_1^j - a > (-x_2^j - a) \tan(\theta)$.

Thus, and since also $z \mapsto \varphi^j(\pm z)$ is analytic in $T_\theta^{\delta, M}$ and continuous in $\overline{T_\theta^{\delta, M}}$, for $0 < \delta < M$, applying Cauchy's integral theorem we have

$$\int_{\partial T_\theta^{\delta, M}} h(x_1^j - a, x_2^j \mp z) \varphi^j(\pm z) dz = 0, \quad \mathbf{x}^j \in \Omega_\theta^j.$$

To complete the proof, we have to show that

$$\lim_{\delta \rightarrow 0} \int_0^\theta h(x_1^j - a, x_2^j \mp (a + \delta e^{i\alpha})) \varphi^j(\pm(a + \delta e^{i\alpha})) i \delta e^{i\alpha} d\alpha = 0$$

and

$$\lim_{M \rightarrow +\infty} \int_0^\theta h(x_1^j - a, x_2^j \mp (a + M e^{i\alpha})) \varphi^j(\pm(a + M e^{i\alpha})) i M e^{i\alpha} d\alpha = 0.$$

These two limits are a consequence of Proposition 3.2, since the constraint $\mathbf{x}^j \in \Omega_\theta^j$ ensures, by (3.17), that $h(x_1^j - a, x_2^j \mp z)$ is a continuous function of z in $\overline{T_\theta}$, and the bound (C.1) and the asymptotics (3.7)–(3.8) imply that $h(x_1^j - a, x_2^j \mp z) = \mathcal{O}(|z|^{-1/2})$ as $z \rightarrow +\infty$ in T_θ , uniformly in $\text{Arg}(z)$. \square

3.3. Derivation and analysis of the complex-scaled HSM method. We derive now an analogue of (2.19) for the complex-scaled traces φ_θ^j . We know, thanks to Theorem 3.4, that

$$u(\mathbf{x}^0) = \int_{\mathbb{R}} h(x_1^0 - a, x_2^0 - \tau_\theta(s)) \varphi_\theta^0(s) \tau'_\theta(s) ds, \quad \mathbf{x}^0 \in \Omega_\theta^0.$$

As $\Omega^0 \cap \Sigma^1 \subset \Omega_\theta^0$, for all $\theta \in (0, \pi/2)$, this holds, in particular, when $\mathbf{x}^0 = (x_1^0, x_2^0) \in \Omega^0 \cap \Sigma^1$, i.e., for $x_1^0 = -x_2^1 > a$ and $x_2^0 = x_1^1 = a$, so that (cf. (2.18))

$$(3.18) \quad \varphi^1(x_2^1) = \int_{\mathbb{R}} h(-x_2^1 - a, a - \tau_\theta(s)) \varphi_\theta^0(s) \tau'_\theta(s) ds, \quad x_2^1 < -a.$$

Remember that $\varphi^1(x_2^1)$ is analytic in $\Re(x_2^1) < -a$ and, by the definition (3.12), $\varphi_\theta^1(t)$ for $t < -a$ is the value of the analytic continuation of φ^1 at $x_2^1 = \tau_\theta(t)$.




FIG. 3.3. The contour $\partial T_{\theta}^{\delta, M}$ and the branch points $z_{\pm} = x_2^j \pm i(x_1^j - a)$ of $h(x_1^j - a, x_2^j - z)$.

To obtain the new complex-scaled compatibility relation, the idea is to complexify x_2^1 in (3.18), i.e., simply to substitute $x_2^1 = \tau_{\theta}(t)$. By the uniqueness of analytic continuation, this is valid provided the right-hand side is analytic as a function of x_2^1 in a connected domain containing the half-line $x_2^1 < -a$ and the half-line $x_2^1 = \tau_{\theta}(t)$, $t < -a$. We check this in the next lemma.

LEMMA 3.6. Let $\psi \in L^2(\mathbb{R})$. For $0 < \theta < \pi/2$ the function

$$(3.19) \quad z \rightarrow \int_{\mathbb{R}} h(z - a, a - \tau_{\theta}(s)) \psi(s) \tau'_{\theta}(s) ds$$

is analytic in the domain $G_{\theta} := \{z \in \mathbb{C} : z \neq a, -\pi/2 + \theta < \text{Arg}(z - a) < \pi/2\}$.

Proof. We will prove that this function is analytic using standard results about analyticity of functions defined as integrals based on the dominated convergence theorem (e.g., [3, Corollary X.3.18]). By the definition (2.14), for each $s \in \mathbb{R}$ the kernel $h(z - a, a - \tau_{\theta}(s))$ is locally an analytic function of z wherever the quantity

$$R(z - a, \tau_{\theta}(s) - a) = [(z - a)^2 + (\tau_{\theta}(s) - a)^2]^{1/2}$$

does not vanish. Lemma B.5 shows that, for every $\theta_0 \in (\theta, \pi/2)$,

$$(3.20) \quad |R(z - a, \tau_{\theta}(s) - a)|^2 \geq \cos(\theta_0) |z - a|^2$$

if $-\theta_0 + \theta \leq \text{Arg}(z - a) \leq \theta_0$. Consequently, $R(z - a, \tau_{\theta}(s) - a)$ does not vanish in G_{θ} .

Further, (3.20) implies that, for every bounded subdomain U of G_{θ} that is bounded away from a , there exists $m_U > 0$ such that

$$|R(z - a, \tau_{\theta}(s) - a)| \geq m_U, \quad z \in U, \quad s \in \mathbb{R}.$$

Moreover, for any such subdomain it follows from Lemma B.4 that, for some constant $C > 0$ independent of $s \in \mathbb{R}$ and $z \in U$,

$$\Im(R(z - a, \tau_{\theta}(s) - a)) \geq |s| \sin(\theta) - C.$$

From the above bounds, and the bound (C.1), it follows that

$$|h(z - a, a - \tau_{\theta}(s))| \leq C' e^{-k \sin(\theta) |s|}$$

for some constant $C' > 0$ independent of $s \in \mathbb{R}$ and $z \in U$. Thus, for every $z \in U$ and $s \in \mathbb{R}$, the integrand in (3.19) has modulus $\leq H(s)$, where $H \in L^1(\mathbb{R})$ is defined by

$$H(s) := C' e^{-k \sin(\theta)|s|} |\psi(s)|, \quad s \in \mathbb{R}.$$

This domination property, and the analyticity of $z \mapsto h(z-a, a-\tau_\theta(s))$ in $U \subset G_\theta$, imply (e.g., [3, Corollary X.3.18]) that the function (3.19) is analytic in U , for every U , and so analytic in G_θ . \square

Noting that, with G_θ as defined in the above lemma, $-x_2^1 \in G_\theta$ for $x_2^1 < -a$ and $-\tau_\theta(t) \in G_\theta$ for $t < -a$, we see that we have justified the analytic continuation of (3.18) from $x_2^1 < -a$ to the path $x_2^1 = \tau_\theta(t)$, $t < -a$. Thus we obtain finally the new complex-scaled compatibility relation

$$\varphi_\theta^1(t) = \int_{\mathbb{R}} h(-\tau_\theta(t) - a, a - \tau_\theta(s)) \varphi_\theta^0(s) \tau_\theta'(s) ds, \quad t < -a.$$

By applying similar reasoning, and noting that $\tau_\theta(-t) = -\tau_\theta(t)$, we get eight equations linking the four complex-scaled traces (cf. (2.19)), namely

$$(3.21) \quad \forall j \in \llbracket 0, 3 \rrbracket, \quad \begin{cases} \varphi_\theta^j(t) = S D_\theta \varphi_\theta^{j-1}(t), & t < -a, \\ \varphi_\theta^j(t) = D_\theta S \varphi_\theta^{j+1}(t), & t > a, \end{cases}$$

where we have set $\varphi_\theta^{-1} := \varphi_\theta^3$ and $\varphi_\theta^4 := \varphi_\theta^0$. In this system the operator S is defined as in (2.20), and, where h is as given in (2.14), D_θ is defined, for $0 < \theta < \pi/2$, by

$$(3.22) \quad D_\theta \psi(t) := \int_{\mathbb{R}} h(\tau_\theta(t) - a, a - \tau_\theta(s)) \psi(s) \tau_\theta'(s) ds, \quad t > a, \quad \psi \in L^2(\mathbb{R}),$$

and, similarly to (2.27), by

$$D_\theta \psi(t) := 0, \quad t \leq a, \quad \psi \in L^2(\mathbb{R}).$$

The system (3.21) has to be completed with the Dirichlet boundary condition

$$(3.23) \quad \varphi_\theta^j(t) = g|_{\Sigma_a^j}(t), \quad -a < t < a, \quad j \in \llbracket 0, 3 \rrbracket.$$

As in section 2, these equations can be formulated as a single operator equation. Introducing

$$\Phi_\theta := \{\varphi_\theta^0, \varphi_\theta^1, \varphi_\theta^2, \varphi_\theta^3\} \in (L^2(\mathbb{R}))^4,$$

and recalling the definition (2.30) of Φ_g , the systems of equations (3.21) and (3.23) can be rewritten as

$$(3.24) \quad \text{Find } \Phi_\theta \in (L^2(\mathbb{R}))^4 \text{ such that } \Phi_\theta - \Phi_g \in (L_0^2(\mathbb{R}))^4 \text{ and} \\ (\mathbb{I} - \mathbb{D}_\theta)(\Phi_\theta - \Phi_g) = \mathbb{D}_\theta \Phi_g,$$

where \mathbb{D}_θ is obtained by replacing D by D_θ in (2.32), i.e.,

$$(3.25) \quad \mathbb{D}_\theta := \begin{bmatrix} 0 & D_\theta S & 0 & S D_\theta \\ S D_\theta & 0 & D_\theta S & 0 \\ 0 & S D_\theta & 0 & D_\theta S \\ D_\theta S & 0 & S D_\theta & 0 \end{bmatrix}.$$

Our first main result is that, as proved for \mathbb{D} in the dissipative case (Theorem 2.2), the operator $\mathbb{I} - \mathbb{D}_\theta$, as an operator on $(L_0^2(\mathbb{R}))^4$, is Fredholm of index zero, indeed (importantly for numerical analysis of Galerkin methods) is a compact perturbation of a coercive operator.

THEOREM 3.7. For $0 < \theta < \pi/2$:

- (i) \mathbb{D}_θ is a continuous operator on $(L^2(\mathbb{R}))^4$, and $\mathbb{D}_\theta((L^2(\mathbb{R}))^4) \subset (L_0^2(\mathbb{R}))^4$;
- (ii) as an operator on $(L_0^2(\mathbb{R}))^4$, \mathbb{D}_θ is the sum of an operator of norm $\leq 1/\sqrt{2}$ and a compact operator.

To show well-posedness of (3.24), it remains to prove a uniqueness result which is the subject of section 5 (Theorem 5.1). At the end of this section we will combine the above theorem with Theorem 5.1 to write down a result expressing this well-posedness and the equivalence of (3.24) with the original scattering problem (3.1)–(1.2).

The proof of Theorem 3.7, which we defer to Appendix A, mirrors the proof of Theorem 2.2, once we establish properties of the operator D_θ to mirror those proved for D in Proposition 2.1. Establishing these properties of D_θ , in Propositions 3.8, 3.9, and 3.10, is the focus of most of the rest of this section: these propositions give the properties of D_θ when it acts on functions whose support is, respectively, in $(a, +\infty)$, $(-\infty, -a)$, and the whole of \mathbb{R} . (This splitting is necessary because of the piecewise definition of the complex-scaling function τ_θ .)

Let us point out a useful fact (see the proof of the following proposition): when D_θ acts on functions whose support is in $(a, +\infty)$, it is equal to the operator D defined in (2.21) for the dissipative case with wavenumber $ke^{i\theta}$.

PROPOSITION 3.8. Suppose that $\theta \in (0, \pi/2)$. For all $\psi \in L^2(a, +\infty)$ we have

$$(3.26) \quad D_\theta \psi(t) = \frac{i(t-a)ke^{i\theta}}{2} \int_a^{+\infty} \frac{H_1^{(1)}(ke^{i\theta} R(t-a, s-a))}{R(t-a, s-a)} \psi(s) ds, \quad t > a,$$

with R defined in (2.15). As a consequence, as an operator on $L^2(a, +\infty)$, D_θ is the sum of an operator of norm $\leq 1/\sqrt{2}$ and a compact operator.

Proof. Using the definition (3.22) of D_θ and the expression (2.14) for the kernel h , we easily see (3.26). This implies that D_θ , when it acts on functions whose support is in $(a, +\infty)$, is exactly the operator D defined in (2.21) for the dissipative case if we set the wavenumber in the dissipative case to be $ke^{i\theta}$. The result is therefore a direct consequence of item (ii) of Proposition 2.1. \square

PROPOSITION 3.9. D_θ is compact as an operator from $L^2(-\infty, -a)$ to $L^2(a, +\infty)$ for $\theta \in (0, \pi/2)$.

Proof. Using the definition (3.22) of D_θ and the expression (2.14) for the kernel h , we can show easily that, for all $\psi \in L^2(-\infty, -a)$, we have

$$D_\theta \psi(t) = \frac{ik(t-a)e^{2i\theta}}{2} \int_{-\infty}^{-a} \frac{H_1^{(1)}(kR_\theta(t, s))}{R_\theta(t, s)} \psi(s) ds, \quad t > a,$$

with $R_\theta(t, s) := R(e^{i\theta}(t-a), 2a-(s+a)e^{i\theta})$ and R defined in (2.15). We prove the compactness of D_θ , as we prove the compactness of D in (iii) of Proposition 2.1, by showing that D_θ is a Hilbert–Schmidt operator from $L^2(-\infty, -a)$ to $L^2(a, +\infty)$. By a simple change of variable $t \mapsto t-a$ and $s \mapsto -(s+a)$, it suffices to show that

$$(3.27) \quad \int_0^{+\infty} \int_0^{+\infty} |K(t, s)|^2 dt ds < +\infty,$$

where

$$K(t, s) := t \frac{H_1^{(1)}(k\tilde{R}_\theta(t, s))}{\tilde{R}_\theta(t, s)} \quad \text{and} \quad \tilde{R}_\theta(t, s) := R(e^{i\theta}t, 2a+e^{i\theta}s).$$

Where $\mathbb{R}^+ := [0, +\infty)$, K is continuous on $\mathbb{R}^+ \times \mathbb{R}^+$, since $\Re(\tilde{R}_\theta) > 0$ on $\mathbb{R}^+ \times \mathbb{R}^+$. We want to use now the bound on the Hankel function given in (C.1) which implies that, for some $c > 0$,

$$(3.28) \quad \left| \frac{H_1^{(1)}(z)}{z} \right| \leq c|z|^{-3/2} e^{-\Im(z)}, \quad \Re(z) > 0, |z| \geq 1.$$

\tilde{R}_θ can be rewritten as $\tilde{R}_\theta(t, s) = \hat{R}(\hat{z}, z)$, where $\hat{z} := 4ase^{i\theta} + 4a^2$, $z := e^{i\theta}(t^2 + s^2)^{1/2}$, and \hat{R} is defined in (B.1). Thus we can use Lemma B.4 and deduce that, for some $C > 0$ and all $s, t \in \mathbb{R}^+$ with $s^2 + t^2$ sufficiently large, it holds that

$$|\tilde{R}_\theta(t, s) - e^{i\theta}(t^2 + s^2)^{1/2}| \leq C \frac{1 + |s|}{(s^2 + t^2)^{1/2}}$$

and

$$\Im(\tilde{R}_\theta(t, s)) \geq \sin(\theta)(t^2 + s^2)^{1/2} - C \frac{1 + |s|}{(s^2 + t^2)^{1/2}}.$$

Thus, for some $C' > 0$ and all $s^2 + t^2$ large enough, $|\tilde{R}_\theta(t, s)| \geq \frac{1}{2}(t^2 + s^2)^{1/2}$ and $\Im(\tilde{R}_\theta(t, s)) \geq \sin(\theta)(t^2 + s^2)^{1/2} - C'$, so that, by (3.28),

$$|K(t, s)| = \mathcal{O} \left(\frac{e^{-k \sin(\theta) \sqrt{t^2 + s^2}}}{(t^2 + s^2)^{1/4}} \right) \quad \text{as } \sqrt{t^2 + s^2} \rightarrow +\infty,$$

uniformly in t and s . Thus (3.27) is clear. \square

PROPOSITION 3.10. For $0 < \theta < \pi/2$, D_θ is a continuous operator from $L^2(\mathbb{R})$ to $L^2(a, +\infty)$.

Proof. From Propositions 3.8 and 3.9, it suffices to show that D_θ is a continuous operator from $L^2(-a, a)$ to $L^2(a, +\infty)$. But this is immediate from the continuity of \mathcal{D}_θ^j from $L^2(\Sigma_a)$ to $L^2(a, +\infty)$, established in Proposition C.4. \square

We finish this section with the promised statement of well-posedness of (3.24), and of its equivalence with the original scattering problem (3.1)–(1.2).

THEOREM 3.11. For every $\theta \in (0, \pi/2)$, the operator $\mathbb{I} - \mathbb{D}_\theta$ is invertible on $(L_0^2(\mathbb{R}))^4$. Thus, for every $g \in L^2(\Sigma_a)$, (3.24) has exactly one solution $\Phi_\theta \in (L^2(\mathbb{R}))^4$ such that $\Phi_\theta - \Phi_g \in (L_0^2(\mathbb{R}))^4$. Moreover, for some constant $c > 0$ depending on θ ,

$$(3.29) \quad \|\Phi_\theta\|_{(L^2(\mathbb{R}))^4} \leq c \|\Phi_g\|_{(L^2(\mathbb{R}))^4} = c \|g\|_{L^2(\Sigma_a)}$$

for every $g \in L^2(\Sigma_a)$. Further, if $g \in H^{1/2}(\Sigma_a)$ and $\Phi_\theta = \{\varphi_\theta^0, \varphi_\theta^1, \varphi_\theta^2, \varphi_\theta^3\} \in (L^2(\mathbb{R}))^4$ is the solution of (3.24), then, for $j \in \llbracket 0, 3 \rrbracket$,

- (i) φ_θ^j satisfies (3.12), i.e., φ_θ^j is the analytic continuation to the path τ_θ of the restriction to Σ^j of the solution u of (3.1)–(1.2);
- (ii) the solution u of (3.1)–(1.2) is given in terms of φ_θ^j in Ω_θ^j by (3.15);
- (iii) for some constant $C > 0$ that depends only on a, k , and θ , $\varphi_\theta^j(s)$ satisfies the bounds of Proposition 3.3 for $|s| > a$.

Proof. Theorem 3.7 implies that, as an operator on $(L_0^2(\mathbb{R}))^4$, $\mathbb{I} - \mathbb{D}_\theta$ is Fredholm of index zero, and Theorem 5.2 implies that it is injective, so that $\mathbb{I} - \mathbb{D}_\theta$ is invertible with a bounded inverse. This implies, since \mathbb{D}_θ is a bounded operator from $(L^2(\mathbb{R}))^4$

to $(L_0^2(\mathbb{R}))^4$ by Theorem 3.7(i), that (3.24) has exactly one solution, and this solution satisfies the bound (3.29).

In the case that $g \in H^{1/2}(\Sigma_a)$, that (i) holds follows from the derivation of (3.24) from (3.1)–(1.2) in section 3.3, and since (3.24) has only one solution; that (ii) holds follows from (i) and Theorem 3.4; that (iii) holds follows from (i) and Proposition 3.3. \square

4. Reconstruction of the solution and far-field formula. Suppose that we have computed the solution Φ_θ to (3.24). Then the solution u of the problem (3.1)–(1.2) can be recovered a posteriori through the representation formulas (3.15). More precisely, as we have observed in Theorem 3.11(ii), it can be reconstructed in the union for $j \in \llbracket 0, 3 \rrbracket$ of the domains Ω_θ^j defined by (3.16). Let us point out that if $\theta < \pi/4$, the union of the Ω_θ^j covers the whole domain Ω , so that the whole solution u can be reconstructed a posteriori.

It is well known (e.g., [21, Lemma 2.5]) that the solution of (3.1)–(1.2) satisfies

$$(4.1) \quad u(\mathbf{x}) = \frac{e^{ikr}}{r^{1/2}} (F(\hat{\mathbf{x}}) + \mathcal{O}(r^{-1})) \quad \text{as } r \rightarrow +\infty,$$

uniformly in $\hat{\mathbf{x}} := \mathbf{x}/r$, where $F \in C^\infty(S^1)$, with S^1 the unit circle, is the *far-field pattern*. By analogy with classical boundary integral methods, one can wonder if this far-field pattern can also be recovered from properties of the φ_θ^j . A partial answer will be given in this section, by deriving far-field formulas in the four directions orthogonal to the edges of the square Ω_a . The proof of this result requires first that we establish some properties of the solution u that can be deduced from the representation formulas (3.15). The far-field behavior that we will establish, indeed all of the results of this section, will be ingredients in the proof of uniqueness for problem (3.24) that will be the focus of the next section.

From (3.15), let us consider the representation formula for any $\psi \in L^2(\mathbb{R})$

$$(4.2) \quad U_\theta(\psi)(\mathbf{x}^0) := \int_{\mathbb{R}} h(x_1^0 - a, x_2^0 - \tau_\theta(s)) \psi(s) \tau'_\theta(s) ds, \quad \mathbf{x}^0 \in \Omega_\theta^0,$$

and the associated integral operator defined by

$$(4.3) \quad \tilde{D}_\theta \psi(t) := U_\theta(\psi)(t, a), \quad t > a.$$

Let us note that in Lemma 3.6 we have shown that, for any data $\psi \in L^2(\mathbb{R})$, the function $t \mapsto \tilde{D}_\theta \psi(t)$ can be continued analytically from $(a, +\infty)$ into the domain of the complex plane $\{z \in \mathbb{C} : z \neq a, -\pi/2 + \theta < \text{Arg}(z - a) < \pi/2\}$ just replacing t by z in (4.3). The following proposition bounds $\tilde{D}_\theta \psi(z)$, in particular near a and for large values of z .

LEMMA 4.1. *There exists a constant $C > 0$, depending only on θ , a , and k , such that, for all $\psi \in L^2(\mathbb{R})$ and $z \in \mathbb{C}$ with $\Re(z) > a$ and $\text{Arg}(z - a) \in [0, \theta]$,*

$$|\tilde{D}_\theta \psi(z)| \leq C(|z - a|^{-1/2} + 1) \exp(-k\Im(z)) \|\psi\|_{L^2(\mathbb{R})}.$$

Proof. Throughout this proof C will denote any positive constant, depending only on a , k , and θ , not necessarily the same at each occurrence.

Using the expressions (4.2)–(4.3), the definition (2.14) of the kernel h , and the definition of the complex-scaling function (3.11), we see that $|\tilde{D}_\theta \psi(z)| \leq \mathcal{I}(z - a)$,

where

$$\mathcal{I}(w) := \frac{k|w|}{2} \int_{\mathbb{R}} \left| \frac{H_1^{(1)}(kR(w, \tau_{\theta}(s) - a))}{R(w, \tau_{\theta}(s) - a)} \right| |\psi(s)| ds, \quad \Re(w) > 0, \quad \text{Arg}(w) \in [0, \theta],$$

with R defined in (2.15). The bound on the Hankel function given in (C.1) implies that

$$\left| \frac{H_1^{(1)}(z)}{z} \right| \leq C \left(|z|^{-2} + |z|^{-3/2} \right) e^{-\Im z}, \quad \Re(z) > 0.$$

Further, Lemma B.5 (applied with $\theta_0 = \theta$) gives, for $\text{Arg}(w) \in [0, \theta]$, that

$$|R(w, \tau_{\theta}(s) - a)|^2 \geq \cos(\theta)(|w|^2 + |\tau_{\theta}(s) - a|^2)$$

and

$$\Im R(w, \tau_{\theta}(s) - a) \geq \cos(\theta - \text{Arg}(w)) \Im(w) - C.$$

Letting $t := |w|$ and $\gamma := \text{Arg}(w)$, we deduce from the above bounds that

$$|\mathcal{I}(w)| e^{k \cos(\theta - \gamma) \Im(w)} \leq C \int_{\mathbb{R}} \left(\frac{t}{t^2 + |\tau_{\theta}(s) - a|^2} + \frac{t}{(t^2 + |\tau_{\theta}(s) - a|^2)^{3/4}} \right) |\psi(s)| ds.$$

Applying the Cauchy-Schwarz inequality, and noticing that $|\tau_{\theta}(s) - a|^2 = (s - a)^2$, for $s \geq -a$, while $|\tau_{\theta}(s) - a|^2 = (s + a)^2 + 4a^2(1 - \cos(\theta)) - 4as \cos(\theta) \geq (s + a)^2$, for $s < -a$, yields

$$|\mathcal{I}(w)| e^{k \cos(\theta - \gamma) \Im(w)} \leq C \left[\int_0^{+\infty} \frac{t^2}{(t^2 + s^2)^2} ds + \int_0^{+\infty} \frac{t^2}{(t^2 + s^2)^{3/2}} ds \right]^{1/2} \|\psi\|_{L^2(\mathbb{R})}.$$

We see, by substituting $s = tp$, that the first and second integrals on the right-hand side of this last inequality are $\leq Ct^{-1}$ and $\leq C$, respectively. Thus we have shown that

$$|\tilde{D}_{\theta} \psi(z)| \leq C(|z - a|^{-1/2} + 1) \exp(-k \cos(\theta - \text{Arg}(z - a)) \Im(z - a)) \|\psi\|_{L^2(\mathbb{R})}$$

for all $\psi \in L^2(\mathbb{R})$ and $z \in \mathbb{C}$ with $\Re(z) > a$ and $\text{Arg}(z - a) \in [0, \theta]$. Now, defining

$$\mathcal{F}(z) := (1 + (z - a)^{-1/2})^{-1} \exp(-ikz) \tilde{D}_{\theta} \psi(z),$$

this last bound implies that, for $\Re(z) > a$ with $0 \leq \text{Arg}(z - a) \leq \theta$, \mathcal{F} is analytic and $|\mathcal{F}(z)| \leq C \|\psi\|_{L^2(\mathbb{R})} B(z)$, where $B(z) := \exp(k \Im(z)(1 - \cos(\theta - \text{Arg}(z - a))))$. Now $B(z) = 1$ when $\text{Arg}(z - a) = 0$ or θ , and $|B(z)| \leq \exp(k|z|)$, for all z with $\Re(z) > a$ and $0 \leq \text{Arg}(z - a) \leq \theta$. Thus, by a standard Phragmén-Lindelöf principle (e.g., [27, Chapter VI, Corollary 4.2]), $|\mathcal{F}(z)| \leq C \|\psi\|_{L^2(\mathbb{R})}$, for all z with $\Re(z) > a$ and $0 \leq \text{Arg}(z - a) \leq \theta$, and the required bound on $|\tilde{D}_{\theta} \psi(z)|$ follows. \square

Defining $\phi(t) := \tilde{D}_{\theta} \psi(t)$, for $t > a$, Lemma 4.1 implies that $\phi \in L^1(a, b)$, for every $b > a$, if $\psi \in L^2(\mathbb{R})$. In the uniqueness proof in the next section we will need also the following stronger result.

LEMMA 4.2. *If $\psi \in L^2(\mathbb{R})$ and $\phi(t) := \tilde{D}_{\theta} \psi(t)$, for $t > a$, then $\phi \in L^2(a, b)$, for every $b > a$.*

Proof. Arguing as in the proof of Lemma 4.1, we see that

$$|\phi(t)| \leq C \left(\int_{\mathbb{R}} \frac{(t-a)|\psi(s)|}{(t-a)^2 + |\tau_{\theta}(s) - a|^2} ds + \int_{\mathbb{R}} \frac{(t-a)|\psi(s)|}{((t-a)^2 + |\tau_{\theta}(s) - a|^2)^{3/4}} ds \right)$$

for $t > a$. Further, arguing as at the end of the proof of Lemma 4.1, using that $|\tau_{\theta}(s) - a|^2 = (s-a)^2$, for $s \geq -a$, while $|\tau_{\theta}(s) - a|^2 \geq (s+a)^2$, for $s < -a$, we see that the second integral in the above sum is bounded on $(a, +\infty)$, and so is in $L^2(a, b)$, for every $b > a$, while the first integral is

$$\leq \int_{-a}^{+\infty} \frac{t-a}{(t-a)^2 + (s-a)^2} |\psi(s)| ds + \int_a^{+\infty} \frac{t-a}{(t-a)^2 + (s-a)^2} |\psi(-s)| ds.$$

The right-hand side of this last inequality is in $L^2(a, +\infty)$ by Remark C.3. \square

Let us remark that by definition (3.22) we have, for any $\psi \in L^2(\mathbb{R})$,

$$(4.4) \quad D_{\theta}\psi(t) = \tilde{D}_{\theta}\psi(\tau_{\theta}(t)) = U_{\theta}(\psi)(\tau_{\theta}(t), a), \quad t > a.$$

We deduce thus from Lemma 4.1 the following result.

COROLLARY 4.3. *There exists a constant $C > 0$, depending only on θ , a , and k , such that, for all $\psi \in L^2(\mathbb{R})$,*

$$|D_{\theta}\psi(t)| \leq C[(t-a)^{-1/2} + 1] e^{-k \sin(\theta)t} \|\psi\|_{L^2(\mathbb{R})}, \quad t > a.$$

Now we are able to prove the main result of this section, which provides far-field formulas in the direction orthogonal to the edges of the square Ω^a .

PROPOSITION 4.4. *Let $\psi \in L^2(\mathbb{R})$ be such that $(t^2 + 1)\psi(t) \in L^1(\mathbb{R})$. Then the function $U_{\theta}(\psi)$ defined by (4.2) has the following behavior at infinity:*

$$U_{\theta}(\psi)(x_1, x_2) = C_{\infty} \frac{e^{ikx_1}}{\sqrt{x_1}} (1 + \mathcal{O}(x_1^{-1})), \quad \text{as } x_1 \rightarrow +\infty, \text{ for each } x_2 \in [-a, a] \text{ fixed,}$$

where

$$C_{\infty} := \sqrt{\frac{k}{\pi}} \frac{1-i}{2} \int_{\mathbb{R}} \psi(s) \tau'_{\theta}(s) ds.$$

Proof. Throughout this proof, C will denote a positive constant, depending only on a , k , and θ , not necessarily the same at each occurrence.

For any fixed $x_2 \in [-a, a]$, let $\phi_{x_2}(t) := U_{\theta}(\psi)(t + a, x_2)$ for $t > 0$. By the definition (2.14) of h , we have

$$\phi_{x_2}(t) = \frac{ikt}{2} \int_{\mathbb{R}} \frac{H_1^{(1)}(kR(t, x_2 - \tau_{\theta}(s)))}{R(t, x_2 - \tau_{\theta}(s))} \psi(s) \tau'_{\theta}(s) ds.$$

In order to use (3.6) that says that

$$(4.5) \quad H_1^{(1)}(z) = -\frac{1+i}{\sqrt{\pi}} \frac{e^{iz}}{\sqrt{z}} \left(1 + \mathcal{O}(|z|^{-1})\right) \quad \text{as } |z| \rightarrow +\infty,$$

uniformly in $\text{Arg}(z)$ for $|\text{Arg}(z)| < \zeta$, for every $\zeta < \pi$, we write

$$(4.6) \quad \phi_{x_2}(t) - \sqrt{\frac{k}{\pi}} \frac{1-i}{2} \frac{e^{ikt}}{\sqrt{t}} \int_{\mathbb{R}} \psi(s) \tau'_{\theta}(s) ds = \phi_{x_2}^1(t) + \phi_{x_2}^2(t),$$

where

$$\begin{aligned}\phi_{x_2}^1(t) &:= \sqrt{\frac{k}{\pi}} \frac{1-i}{2} t \int_{\mathbb{R}} \left[\frac{e^{ikR(t, x_2 - \tau_\theta(s))}}{[R(t, x_2 - \tau_\theta(s))]^{3/2}} - \frac{e^{ikt}}{t^{3/2}} \right] \psi(s) \tau'_\theta(s) ds, \\ \phi_{x_2}^2(t) &:= \frac{ikt}{2} \int_{\mathbb{R}} \left[\frac{H_1^{(1)}(kR(t, x_2 - \tau_\theta(s)))}{R(t, x_2 - \tau_\theta(s))} \right. \\ &\quad \left. + \frac{1+i}{\sqrt{k\pi}} \frac{e^{ikR(t, x_2 - \tau_\theta(s))}}{[R(t, x_2 - \tau_\theta(s))]^{3/2}} \right] \psi(s) \tau'_\theta(s) ds.\end{aligned}$$

To estimate these quantities we note first that if $t > 0$ and $z \in Z_\theta := \{z = re^{i\alpha} : 0 \leq \alpha \leq \theta, r \geq 0\}$ (or $-z \in Z_\theta$), then $0 \leq \text{Arg}(R(t, z)) \leq \pi/2$, $|\cos(\text{Arg}(z))| \geq \cos(\theta)$, and, by Lemma B.1,

$$(4.7) \quad |R(t, z)| \geq \cos(\theta) t.$$

Since, from the definition (3.11), $x_2 - \tau_\theta(s)$ or $\tau_\theta(s) - x_2$ is in Z_θ for all $x_2 \in [-a, a]$ and all $s \in \mathbb{R}$, these observations hold in particular if $z = x_2 - \tau_\theta(s)$, so that (4.5) applies and implies that

$$(4.8) \quad |\phi_{x_2}^2(t)| \leq \frac{C}{t^{3/2}} \int_{\mathbb{R}} |\psi(s)| ds$$

for all sufficiently large $t > 0$. To estimate $\phi_{x_2}^1(t)$ we observe that, for $t > 0$ and $z \in Z_\theta$,

$$\left| \frac{e^{ikR(t, z)}}{[R(t, z)]^{3/2}} - \frac{e^{ikt}}{t^{3/2}} \right| \leq \sup_{w \in U(z)} \left| \frac{\partial}{\partial w} \left[\frac{e^{ikR(t, w)}}{[R(t, w)]^{3/2}} \right] \right| |z|,$$

where $U(z) := \{w \in Z_\theta : |w| \leq |z|\}$ and where we have used that $R(t, 0) = t$. Using again (4.7), this yields that

$$\left| \frac{e^{ikR(t, z)}}{[R(t, z)]^{3/2}} - \frac{e^{ikt}}{t^{3/2}} \right| \leq C \frac{|z|^2}{t^{5/2}}, \quad z \in Z_\theta, \quad t > 0.$$

Since $|x_2 - \tau_\theta(s)| \leq C(1 + |s|)$, it follows that

$$(4.9) \quad |\phi_{x_2}^1(t)| \leq \frac{C}{t^{3/2}} \int_{\mathbb{R}} (1 + |s|)^2 |\psi(s)| ds, \quad t > 0.$$

Combining (4.6), (4.8), and (4.9), we get that

$$\left| \phi_{x_2}(t) - \sqrt{\frac{k}{\pi}} \frac{1-i}{2} \frac{e^{ikt}}{\sqrt{t}} \int_{\mathbb{R}} \psi(s) \tau'_\theta(s) ds \right| \leq \frac{C}{t^{3/2}} \int_{\mathbb{R}} (1 + s^2) |\psi(s)| ds$$

for all sufficiently large $t > 0$, which ends the proof. \square

This proposition proves in particular that

$$(4.10) \quad F(\cos(j\pi/2), \sin(j\pi/2)) = \sqrt{\frac{k}{\pi}} \frac{1-i}{2} \int_{\mathbb{R}} \varphi_\theta^j(s) \tau'_\theta(s) ds, \quad j \in \llbracket 0, 3 \rrbracket,$$

where the far-field F is defined by (4.1) and $\varphi_\theta^0, \varphi_\theta^1, \varphi_\theta^2, \varphi_\theta^3$ are the complex-scaled traces of u , which are exponentially decaying at infinity thanks to Proposition 3.3, or thanks to (3.21) and Corollary 4.3.

5. Uniqueness. In this section we will prove uniqueness of solution for the complex-scaled HSM method (3.24). This result, important in its own right, is also key to the proof of uniqueness for the complex-scaled HSM method for more general configurations; see Proposition 6.1 below. Our proof depends on the following uniqueness result for the standard HSM (2.25) that we prove, using completely different arguments, in [9].

THEOREM 5.1. *If $k > 0$ and $\{\varphi^0, \varphi^1, \varphi^2, \varphi^3\} \in (L^2_{\text{loc}}(\mathbb{R}))^4$ satisfies (2.25) with $g = 0$, and satisfies the radiation condition that*

$$(5.1) \quad \varphi^j(t) = \begin{cases} c_+^j e^{ik|t|} |t|^{-1/2} (1 + \mathcal{O}(|t|^{-1})) & \text{as } t \rightarrow +\infty, \\ c_-^j e^{ik|t|} |t|^{-1/2} (1 + \mathcal{O}(|t|^{-1})) & \text{as } t \rightarrow -\infty, \end{cases}$$

for some constants $c_{\pm}^j \in \mathbb{C}$ and every $j \in \llbracket 0, 3 \rrbracket$, then $\varphi^j = 0$ for $j \in \llbracket 0, 3 \rrbracket$.

To see how this result is relevant to uniqueness for the complex-scaled HSM method (3.24), recall that to derive (3.21) and (3.23) we started from (2.25), satisfied for real k by φ^j , the trace on Σ^j of the solution u of (3.1)–(1.2), for $j \in \llbracket 0, 3 \rrbracket$. We showed that φ_{θ}^j , defined by (3.12) as the analytic continuation of φ^j from the real line to the path τ_{θ} of Figure 3.1, satisfies (3.21) and (3.23), equivalently (3.24). A key component in this argument was to deform paths of integration from the real line to the path τ_{θ} (Theorem 3.4).

In the following uniqueness proof we reverse this derivation. We show that if the φ_{θ}^j satisfy (3.21) and (3.23), with $g = 0$, then, in the sense that (3.12) holds, they are the analytic continuations onto the path τ_{θ} of functions φ^j that satisfy the system (2.25) with $g = 0$. (A key component in this argument is a deformation of paths of integration from τ_{θ} back to the real line, justified as in the proof of Theorem 3.4.) Moreover, by an application of Proposition 4.4 (justified by Corollary 4.3), the functions φ^j satisfy the radiation conditions (5.1), so that $\varphi^j = 0$ by Theorem 5.1. Thus φ_{θ}^j , which is the analytic continuation of φ^j , is also zero.

THEOREM 5.2. *Suppose that $\{\varphi_{\theta}^0, \varphi_{\theta}^1, \varphi_{\theta}^2, \varphi_{\theta}^3\} \in (L^2(\mathbb{R}))^4$ is a solution of (3.21) such that*

$$(5.2) \quad \varphi_{\theta}^j(t) = 0, \quad -a < t < a, \quad j \in \llbracket 0, 3 \rrbracket.$$

Then $\varphi_{\theta}^j = 0$ for $j \in \llbracket 0, 3 \rrbracket$.

Proof. Suppose that $\{\varphi_{\theta}^0, \varphi_{\theta}^1, \varphi_{\theta}^2, \varphi_{\theta}^3\} \in (L^2(\mathbb{R}))^4$ is a solution of (3.21) that satisfies (5.2), and define the functions $\varphi^0, \varphi^1, \varphi^2, \varphi^3$ by

$$(5.3) \quad \begin{cases} \varphi^j(t) := S \tilde{D}_{\theta} \varphi_{\theta}^{j-1}(t), & t < -a, \\ \varphi^j(t) := 0, & -a \leq t \leq a, \\ \varphi^j(t) := \tilde{D}_{\theta} S \varphi_{\theta}^{j+1}(t), & t > a, \end{cases}$$

for $j \in \llbracket 0, 3 \rrbracket$, where \tilde{D}_{θ} is defined by (4.3). As discussed below (4.3), since the φ_{θ}^j 's are in $L^2(\mathbb{R})$, it follows from this definition, the definition (2.20) of S , and Lemma 3.6 that the φ^j 's have analytic continuations from $(a, +\infty)$ (respectively, $(-\infty, -a)$) to the part of the half-plane $\Re(z) > a$ with $-\pi/2 + \theta < \text{Arg}(z - a) < \pi/2$ (respectively, $\Re(z) < -a$ with $-\pi/2 + \theta < \text{Arg}(-z - a) < \pi/2$). In particular, by (4.4), and recalling

(2.20) and that $\tau_\theta(-t) = \tau_\theta(t)$ for $t < -a$,

$$\begin{cases} \varphi^j(\tau_\theta(t)) = S D_\theta \varphi_\theta^{j-1}(t), & t < -a, \\ \varphi^j(\tau_\theta(t)) = 0, & -a \leq t \leq a, \\ \varphi^j(\tau_\theta(t)) = D_\theta S \varphi_\theta^{j+1}(t), & t > a, \end{cases}$$

for $j \in \llbracket 0, 3 \rrbracket$. Comparing this equation with (3.21) we see that

$$(5.4) \quad \varphi_\theta^j(s) = \varphi^j(\tau_\theta(s)) \quad \text{for } s \in \mathbb{R} \text{ and } j \in \llbracket 0, 3 \rrbracket.$$

Thus, for $s > a$ and for $s < -a$, the φ_θ^j 's are the analytic continuations of the φ^j 's to the complex path parametrized by τ_θ . Therefore, to complete the proof of the theorem it is enough to show, for $j \in \llbracket 0, 3 \rrbracket$, that $\varphi^j(t) = 0$ for $t \in \mathbb{R}$ with $|t| > a$, for this will imply, by the uniqueness of analytic continuation, that $\varphi^j(\tau_\theta(s)) = 0$ for $s > a$ and $s < -a$, which will imply, using (5.4), that each $\varphi_\theta^j = 0$.

To establish, for $j \in \llbracket 0, 3 \rrbracket$, that $\varphi^j(t) = 0$ for $t > a$ and $t < -a$, we show, via an application of Cauchy's integral theorem as in the proof of Theorem 3.4, that uses the analyticity of the φ^j 's noted above, Lemma 4.1, and (5.4), that (5.3) implies that the φ^j satisfy (2.25) with $g = 0$, i.e.,

$$\begin{cases} \varphi^j(t) = S D \varphi^{j-1}(t), & t < -a, \\ \varphi^j(t) = 0, & -a < t < a, \\ \varphi^j(t) = D S \varphi^{j+1}(t), & t > a, \end{cases}$$

for $j \in \llbracket 0, 3 \rrbracket$, where D is defined in (2.23), or equivalently by (3.22) with $\theta = 0$. (The key step in the argument is to show, as we have done in the proof of Theorem 3.4, that

$$\int_a^{+\infty} h(t-a, a \mp \tau_\theta(s)) \varphi(s) \tau'_\theta(s) ds = \int_a^{+\infty} h(t-a, a \mp s) \varphi(s) ds$$

if, for every $0 < \delta < M$, φ is analytic in the domain $T_\theta^{\delta, M}$ introduced in the proof of Theorem 3.4, and continuous in its closure, and if the behavior of $\varphi(z)$ as $z \rightarrow 0$ and as $z \rightarrow +\infty$ is suitably constrained, satisfying the bounds of Lemma 4.1.) We note also that it follows from (5.3) and Lemma 4.2 that $\varphi^j \in L^2_{\text{loc}}(\mathbb{R})$ for $j \in \llbracket 0, 3 \rrbracket$. Moreover, the φ_θ^j are in $L^2(\mathbb{R})$ and it follows, from (3.21) and Corollary 4.3, that each φ_θ^j decays exponentially at infinity, so that $(t^2 + 1)\varphi_\theta^j(t) \in L^1(\mathbb{R})$, $j \in \llbracket 0, 3 \rrbracket$. Thus we can apply Proposition 4.4 to (5.3), noting the second equality in (4.4), to see that the radiation conditions (5.1) hold. But this is enough to conclude that $\varphi^j = 0$, for $j \in \llbracket 0, 3 \rrbracket$, by Theorem 5.1. \square

6. The complex-scaled HSM method for the general case. Let us now explain how to extend the complex-scaled HSM method of section 3 to solve the general problem presented in the introduction. More precisely, for a real wavenumber $k > 0$ and for a function ρ and a subdomain Ω of \mathbb{R}^2 satisfying the hypotheses described in section 1, the objective is to derive a complex-scaled HSM formulation to compute the solution $u \in H^1_{\text{loc}}(\Omega)$ of (1.1) and (1.2). For the sake of simplicity, we will restrict attention to the case $\Omega = \mathbb{R}^2$, but adding bounded obstacles contained in Ω_a is completely straightforward (see section 7.3). As in section 1, the source term f is in $L^2(\Omega)$, with compact support that is a subset of Ω_a .

The idea is to introduce, in addition to the lines Σ^j , $j \in \llbracket 0, 3 \rrbracket$, a square $\Omega_b := (-b, b)^2$ for some $b > a$. (As in section 2 we set $\Sigma_b := \partial\Omega_b$ and denote the sides of Σ_b




FIG. 6.1. The notations for the general case.

by Σ_b^j , $j \in \llbracket 0, 3 \rrbracket$; see Figure 6.1.) We will show how to derive a formulation of problem (1.1), (1.2) whose unknowns are the complex-scaled traces φ_θ^j , $j \in \llbracket 0, 3 \rrbracket$, associated to the infinite lines Σ^j , and the restriction $u_b := u|_{\Omega_b}$ of the solution u to the square Ω_b . To do that, we need to make the following assumption on the parameter θ :

$$(6.1) \quad \theta < \frac{\pi}{4}.$$

Let us derive the equations linking the φ_θ^j , $j \in \llbracket 0, 3 \rrbracket$, and u_b . On the one hand, the φ_θ^j still satisfy the system of compatibility relations (3.21). But, instead of (3.23), we have to impose equality between φ_θ^j and u_b on Σ_a^j :

$$\varphi_\theta^j(t) = u_b|_{\Sigma_a^j}(t), \quad -a < t < a, \quad j \in \llbracket 0, 3 \rrbracket.$$

On the other hand, we can derive a variational formulation for u_b in Ω_b . Since $-\Delta u_b - k^2 \rho u_b = f$ in Ω_b and f is supported in Ω_a , the following Green's identity holds for all $v_b \in H^1(\Omega_b)$, where n is the normal unit vector pointing out of Ω_b :

$$(6.2) \quad \int_{\Omega_b} (\nabla u_b \cdot \overline{\nabla v_b} - k^2 \rho u_b \overline{v_b}) - \int_{\Sigma_b} \frac{\partial u_b}{\partial n} \overline{v_b} = \int_{\Omega_a} f \overline{v_b}.$$

The last idea is to replace in the previous identity the normal derivative of u_b on the j th side of the square by an integral representation as a function of φ_θ^j . Indeed, we must have, for $j \in \llbracket 0, 3 \rrbracket$,

$$\frac{\partial u_b}{\partial n} - i k u_b = \frac{\partial U^j}{\partial n} - i k U^j \text{ on } \Sigma_b^j,$$

where U^j denotes the restriction of the solution u to the half-plane Ω^j . (Our choice of Robin traces instead of normal derivatives is so that later we have uniqueness for the boundary value problem (6.8) for all $k > 0$.)

We have proved in Theorem 3.4 that $U^j(\mathbf{x}^j)$ has an integral representation in terms of φ_θ^j , as soon as \mathbf{x}^j belongs to the domain Ω_θ^j defined by (3.16). Notice that under condition (6.1) one has $\Sigma_b^j \subset \Omega_\theta^j$ (see Figure 6.1). Consequently, one can use the formula (3.15) to rewrite the above Robin compatibility condition on Σ_b^j . Precisely,

we have

$$\begin{aligned} & \left(\frac{\partial u_b}{\partial n} - ik u_b \right) \Big|_{\Sigma_b^j}(t) \\ &= \int_{\mathbb{R}} (\partial_1 h(b-a, t - \tau_\theta(s)) - ik h(b-a, t - \tau_\theta(s))) \varphi_\theta^j(s) \tau_\theta'(s) ds, \quad -b < t < b, \end{aligned}$$

where $\partial_1 h$ denotes the derivative of h with respect to its first variable. This leads us to define the following Dirichlet-to-Robin operator Λ_θ . For $\psi \in L^2(\mathbb{R})$ and $0 < \theta < \pi/4$, define $\Lambda_\theta \psi \in L^2(-b, b)$ by

$$(6.3) \quad \Lambda_\theta \psi(t) := \int_{\mathbb{R}} \lambda(b-a, t - \tau_\theta(s)) \psi(s) \tau_\theta'(s) ds, \quad -b < t < b,$$

where we have set

$$\lambda(x_1, z) := \partial_1 h(x_1, z) - ik h(x_1, z), \quad x_1 > 0, \quad z \in \mathbb{C},$$

which one can easily check, using [30, (10.6.2)] and the definition (2.14) of h , takes the explicit form

$$(6.4) \quad \lambda(x_1, z) = \frac{ik}{2R} \left([1 - ikx_1] H_1^{(1)}(kR) - \frac{kx_1^2}{R} H_2^{(1)}(kR) \right), \quad R = [x_1^2 + z^2]^{1/2}.$$

With this notation, the previous equations linking u_b and the φ_θ^j can be written as

$$(6.5) \quad \left(\frac{\partial u_b}{\partial n} - ik u_b \right) \Big|_{\Sigma_b^j}(t) = \Lambda_\theta \varphi_\theta^j(t), \quad -b < t < b, \quad j \in \llbracket 0, 3 \rrbracket.$$

Our complete formulation reads as follows:

$$\begin{aligned} & \text{Find } \{\varphi_\theta^0, \varphi_\theta^1, \varphi_\theta^2, \varphi_\theta^3\} \in (L^2(\mathbb{R}))^4 \text{ and } u_b \in H^1(\Omega_b) \text{ such that} \\ & \quad \begin{cases} \varphi_\theta^j(t) = S D_\theta \varphi_\theta^{j-1}(t) \text{ for } t < -a, \\ \varphi_\theta^j(t) = u_b|_{\Sigma_a^j}(t) \text{ for } -a < t < a, \\ \varphi_\theta^j(t) = D_\theta S \varphi_\theta^{j+1}(t) \text{ for } t > a, \end{cases} \quad j \in \llbracket 0, 3 \rrbracket, \\ (6.6) \quad & \text{and such that, } \forall v_b \in H^1(\Omega_b), \end{aligned}$$

$$\begin{aligned} & \int_{\Omega_b} (\nabla u_b \cdot \overline{\nabla v_b} - k^2 \rho u_b \overline{v_b}) - ik \sum_{j=0}^3 \int_{\Sigma_b^j} u_b \overline{v_b} \\ & - \sum_{j=0}^3 \int_{-b}^b \Lambda_\theta \varphi_\theta^j(t) \overline{v_b}|_{\Sigma_b^j}(t) dt = \int_{\Omega_a} f \overline{v_b}, \end{aligned}$$

where S , D_θ , and Λ_θ are defined by (2.20), (3.22), and (6.3).

Let us denote by Φ_θ , $\Phi(u_b)$, and $\tilde{\Phi}_\theta$ the following elements of $(L^2(\mathbb{R}))^4$:

$$\begin{aligned} \Phi_\theta &:= \{\varphi_\theta^0, \varphi_\theta^1, \varphi_\theta^2, \varphi_\theta^3\}, \quad \Phi(u_b) := \{u_b|_{\Sigma_a^0}, u_b|_{\Sigma_a^1}, u_b|_{\Sigma_a^2}, u_b|_{\Sigma_a^3}\}, \\ \tilde{\Phi}_\theta &:= \{\tilde{\varphi}_\theta^0, \tilde{\varphi}_\theta^1, \tilde{\varphi}_\theta^2, \tilde{\varphi}_\theta^3\} := \Phi_\theta - \Phi(u_b). \end{aligned}$$

Note that, using the second equation of (6.6) and recalling (2.29), we have $\tilde{\Phi}_\theta \in (L_0^2(\mathbb{R}))^4$. With these notations the first block of equations in (6.6) can be rewritten

(cf. (3.24)) as $(\mathbb{I} - \mathbb{D}_\theta)\tilde{\Phi}_\theta = \mathbb{D}_\theta\Phi(u_b)$, where \mathbb{D}_θ is defined by (3.25), so that problem (6.6) can be rewritten as

$$(6.7) \quad \begin{aligned} & \text{Find } \tilde{\Phi}_\theta \in (L_0^2(\mathbb{R}))^4 \text{ and } u_b \in H^1(\Omega_b) \text{ such that} \\ & \quad \forall \tilde{\Psi} \in (L_0^2(\mathbb{R}))^4 \text{ and } \forall v_b \in H^1(\Omega_b), \\ & \quad ((\mathbb{I} - \mathbb{D}_\theta)\tilde{\Phi}_\theta, \tilde{\Psi})_{(L_0^2(\mathbb{R}))^4} - (\mathbb{D}_\theta\Phi(u_b), \tilde{\Psi})_{(L_0^2(\mathbb{R}))^4} \\ & \quad + \int_{\Omega_b} (\nabla u_b \cdot \overline{\nabla v_b} - k^2 \rho u_b \overline{v_b}) - ik \sum_{j=0}^3 \int_{\Sigma_b^j} u_b \overline{v_b} \\ & \quad - \sum_{j=0}^3 \int_{-b}^b \Lambda_\theta \left(\tilde{\varphi}_\theta^j + u_b|_{\Sigma_a^j} \right) (t) \overline{v_b}|_{\Sigma_b^j}(t) dt = \int_{\Omega_a} f \overline{v_b}. \end{aligned}$$

Let us first prove a uniqueness result for this problem.

PROPOSITION 6.1. *If $f = 0$, then the only solution of problem (6.7) is the trivial solution $u_b = 0$, $\tilde{\Phi}_\theta = 0$.*

Proof. Suppose that $\tilde{\Phi}_\theta \in (L_0^2(\mathbb{R}))^4$ and $u_b \in H^1(\Omega_b)$ are such that (6.7) holds with $f = 0$. Then $(\mathbb{I} - \mathbb{D}_\theta)\tilde{\Phi}_\theta = \mathbb{D}_\theta\Phi(u_b)$, and the second of equations (6.6) holds with $f = 0$ and $\varphi_\theta^j := \tilde{\varphi}_\theta^j + u_b|_{\Sigma_a^j}$, $j \in \llbracket 0, 3 \rrbracket$, so that $-\Delta u_b - k^2 \rho u_b = 0$ in Ω_b and (6.5) holds for $j \in \llbracket 0, 3 \rrbracket$.

Let us denote by $u_\infty \in H_{\text{loc}}^1(\mathbb{R}^2 \setminus \overline{\Omega_a})$ the unique solution of (3.1)–(1.2) with $g = u_b|_{\Sigma_a} \in H^{1/2}(\Sigma_a)$. Then, as we have shown in section 3.3, the vector of complex-scaled traces of u_∞ satisfies (3.24) with $\Phi_g = \Phi(u_b)$, so that, by Theorem 5.2, it coincides with the vector $\Phi_\theta := \{\varphi_\theta^0, \varphi_\theta^1, \varphi_\theta^2, \varphi_\theta^3\}$. Thus, applying Theorem 3.4 (as we did above to derive (6.5) from (1.1)–(1.2)) we see that

$$\Lambda_\theta \varphi_\theta^j = \left(\frac{\partial u_\infty}{\partial n} - ik u_\infty \right) \Big|_{\Sigma_b^j}, \quad j \in \llbracket 0, 3 \rrbracket.$$

Consequently, $v := u_b - u_\infty$ belongs to $H^1(\Omega_b \setminus \overline{\Omega_a})$ and satisfies

$$(6.8) \quad \begin{cases} \Delta v + k^2 v = 0 & \text{in } \Omega_b \setminus \overline{\Omega_a}, \\ v = 0 & \text{on } \Sigma_a, \\ \frac{\partial v}{\partial n} - ikv = 0 & \text{on } \Sigma_b. \end{cases}$$

But, for every $k > 0$, this homogeneous problem has no solution except $v = 0$. (To see this apply Green's identity (cf. (6.2)) in $\Omega_b \setminus \overline{\Omega_a}$ to deduce that $\int_{\Sigma_b} |v|^2 = 0$, so that $v = \partial v / \partial n = 0$ on Σ_b , and apply Holmgren's uniqueness theorem; [21, p. 104].) Thus $u_b = u_\infty$ in $\Omega_b \setminus \overline{\Omega_a}$ so that the function

$$w := \begin{cases} u_b & \text{in } \Omega_b, \\ u_\infty & \text{in } \mathbb{R}^2 \setminus \overline{\Omega_a}, \end{cases}$$

is well-defined and is a solution of the homogeneous Helmholtz equation in \mathbb{R}^2 which satisfies the radiation condition (1.2). As a consequence (e.g., [25, Theorem 8.7]), $w = 0$ in \mathbb{R}^2 , so that $u_b = 0$ in Ω_b and $u_\infty = 0$. Further, each φ_θ^j is zero, since it is a complex trace of u_∞ . This completes the proof. \square

The well-posedness of problem (6.7) follows from classical arguments combined with the results of the previous sections and the following lemma.

LEMMA 6.2. *The operator Λ_θ , defined by (6.3), is a compact operator from $L^2(\mathbb{R})$ to $L^2(-b, b)$.*

Proof. To establish this lemma, we will prove that the kernel of Λ_θ is Hilbert-Schmidt, i.e., that

$$(6.9) \quad (t, s) \mapsto \lambda(b-a, t-\tau_\theta(s)) \in L^2((-b, b) \times \mathbb{R}),$$

by using similar arguments as in the proof of Proposition 3.9. On the one hand, since $b > a$ and $\theta < \pi/4$, the function R which appears in expression (6.4) never vanishes for $-b \leq t \leq b$ and $s \in \mathbb{R}$. As a consequence, the kernel is continuous on $[-b, b] \times \mathbb{R}$. On the other hand we have, using (3.7), (3.8), and (3.6), the asymptotic estimate

$$|\lambda(b-a, t-\tau_\theta(s))| = \mathcal{O}\left(\frac{e^{-k \sin \theta |s|}}{|s|^{1/2}}\right) \quad \text{as } |s| \rightarrow +\infty,$$

uniformly in t , for $|t| \leq b$. Together, these properties prove (6.9). \square

Recall that we call a sesquilinear form $a(\cdot, \cdot)$ on a Hilbert space \mathcal{H} *compact* if the associated linear operator A on \mathcal{H} , defined by $(A\phi, \psi) = a(\phi, \psi)$, for all $\phi, \psi \in \mathcal{H}$, is compact. (Here (\cdot, \cdot) denotes the inner product on \mathcal{H} .) Equivalently, $a(\cdot, \cdot)$ is compact if, whenever $\phi_n \rightharpoonup 0$ and $\psi_n \rightharpoonup 0$ (weak convergence in \mathcal{H}), it holds that $a(\phi_n, \psi_n) \rightarrow 0$.

In our final theorem we show that the sesquilinear form on the Hilbert space $(L_0^2(\mathbb{R}))^4 \times H^1(\Omega_b)$ which appears on the left-hand side of (6.7) is the sum of coercive plus compact sesquilinear forms, and that, as a consequence of this and of the above uniqueness result, problem (6.7) is well-posed. Regarding the last sentence of the theorem, note that, given the constraint $0 < \theta < \pi/4$,

$$\Omega \subset \Omega_b \cup \bigcup_{j=0}^3 \Omega_\theta^j,$$

so that the solution of the original scattering problem can be recovered in the whole of Ω from the solution $(\tilde{\Phi}_\theta, u_b)$ of (6.7).

THEOREM 6.3. *For every $\theta \in (0, \pi/4)$ and every $f \in L^2(\Omega)$ with support in Ω_a , problem (6.7) has exactly one solution $(\tilde{\Phi}_\theta, u_b) \in (L_0^2(\mathbb{R}))^4 \times H^1(\Omega_b)$. Further, for some constant $c > 0$ depending on θ ,*

$$(6.10) \quad \|\tilde{\Phi}_\theta\|_{(L_0^2(\mathbb{R}))^4} + \|u_b\|_{H^1(\Omega_b)} \leq c \|f\|_{L^2(\Omega_a)}$$

for all $f \in L^2(\Omega)$ with support in Ω_a . Moreover, if $(\tilde{\Phi}_\theta, u_b) \in (L_0^2(\mathbb{R}))^4 \times H^1(\Omega_b)$ is the solution of (6.7) and $u \in H_{\text{loc}}^1(\Omega)$ is the solution of (1.1)–(1.2), then $u = u_b$ in Ω_b , while u is given in terms of φ_θ^j in Ω_θ^j by (3.15) for $j \in \llbracket 0, 3 \rrbracket$.

Proof. As an operator on $(L_0^2(\mathbb{R}))^4$ we have, by Theorem 3.7, that $\mathbb{D}_\theta = \mathbb{D}_\theta^1 + \mathbb{D}_\theta^2$, where $\|\mathbb{D}_\theta^1\| \leq 1/\sqrt{2}$ and \mathbb{D}_θ^2 is compact. Thus we can decompose the sesquilinear form which appears on the left-hand side of (6.7) as the sum of a first sesquilinear form

$$((\mathbb{I} - \mathbb{D}_\theta^1)\tilde{\Phi}_\theta, \tilde{\Psi})_{(L_0^2(\mathbb{R}))^4} + \int_{\Omega_b} (\nabla u_b \cdot \overline{\nabla v_b} + u_b \overline{v_b}) - ik \sum_{j=0}^3 \int_{\Sigma_b^j} u_b \overline{v_b},$$

which is coercive on $(L_0^2(\mathbb{R}))^4 \times H^1(\Omega_b)$, and a second sesquilinear form

$$\begin{aligned} & -(\mathbb{D}_\theta^2 \tilde{\Phi}_\theta, \tilde{\Psi})_{(L_0^2(\mathbb{R}))^4} - (\mathbb{D}_\theta \Phi(u_b), \tilde{\Psi})_{(L_0^2(\mathbb{R}))^4} - (1 + k^2) \int_{\Omega_b} u_b \overline{v_b} \\ & - \sum_{j=0}^3 \int_{-b}^b \Lambda_\theta \left(\tilde{\varphi}_\theta^j + u_b|_{\Sigma_b^j} \right) (t) \overline{v_b|_{\Sigma_b^j}}(t) dt, \end{aligned}$$

which is compact on the same space. The proofs of compactness of the four terms of this second sesquilinear form rely on different arguments. The first is compact because \mathbb{D}_θ^2 is compact. For the second term, we notice that the operator $u_b \mapsto \mathbb{D}_\theta \Phi(u_b)$ is compact from $H^1(\Omega_b)$ to $(L_0^2(\mathbb{R}))^4$, as it is the composition of the bounded operator \mathbb{D}_θ on $(L_0^2(\mathbb{R}))^4$ and the compact map $u_b \mapsto \Phi(u_b)$. (To see the compactness of this last map, note that it can be thought of as a composition of the bounded trace map from $H^1(\Omega_b)$ to $H^{1/2}(\Sigma_b)$ and the compact embedding $H^{1/2}(\Sigma_b) \subset L^2(\Sigma_b)$.) The compactness of the third term is a consequence of the Rellich embedding theorem, that the embedding $H^1(\Omega_b) \subset L^2(\Omega_b)$ is compact. To see that the last term is compact one can use Lemma 6.2. Indeed, note that continuity of Λ_θ , combined with compactness of the map $v_b \mapsto v_b|_{\Sigma_b^j}$ from $H^1(\Omega_b)$ to $L^2(\Sigma_b^j)$, suffices to conclude.

Since the sesquilinear form is coercive plus compact, the Fredholm alternative holds (e.g., [40, Theorem 2.33]), so that unique solvability of (6.7) and the stability bound (6.10) are a consequence of Proposition 6.1. We have shown the last sentence of the theorem in our derivation, earlier in this section, of (6.7) from (1.1)–(1.2), using Theorem 3.4. \square

7. Numerical implementation and results. In this section we demonstrate, through some illustrative numerical experiments implemented in XLiFE++ [38], that the complex-scaled HSM formulations (3.24) and (6.7) can be solved numerically to compute solutions to the scattering problems (3.1)–(1.2) and (1.1)–(1.2), respectively.

7.1. Numerical implementation of the deformed half-space representation. Before considering the discretization of the HSM systems, we just want to provide an illustration of Theorem 3.4. More precisely, let

$$(7.1) \quad u(\mathbf{x}) := \frac{i}{4} H_0^{(1)}(k R(x_1, x_2)), \quad \mathbf{x} \in \mathbb{R}^2 \setminus \{0\},$$

where R is defined by (2.15), so that u satisfies (3.1)–(1.2) in the case that $g := u|_{\Sigma_a}$. Then $\varphi^0 := u|_{\Sigma^0}$ is given by

$$\varphi^0(s) = \frac{i}{4} H_0^{(1)}(k R(s, a)), \quad s \in \mathbb{R}.$$

The corresponding complex-scaled trace φ_θ^0 , defined by (3.11) and (3.12), is an even function given, for $|s| < a$, by

$$\varphi_\theta^0(s) := \varphi^0(s)$$

and, for $s > a$, by

$$(7.2) \quad \begin{aligned} \varphi_\theta^0(s) := \varphi^0(\tau_\theta(s)) &= \frac{i}{4} H_0^{(1)} \left(k \sqrt{a^2 + (a + (s - a)e^{i\theta})^2} \right) \\ &\sim \frac{e^{i(ka + \pi/4 - \theta/2)} e^{ik(s-a)e^{i\theta}}}{2\sqrt{2\pi ks}} \quad \text{as } s \rightarrow +\infty, \end{aligned}$$




FIG. 7.1. Left: representation of the exact complex-scaled traces $s \mapsto \varphi_\theta^0(s)$ for $\theta = 0, \pi/6, \pi/4, \pi/3$. Right: comparison of the exact (red line) and computed (blue dots) complex-scaled trace $s \mapsto \varphi_\theta^0(s)$ for $\theta = \pi/6$. In both plots $a = 1$ and $k = 2\pi$.

by (3.6) and (3.7). The asymptotic behavior (7.2) agrees with (3.13) and Proposition 3.3, indeed demonstrates that (3.13) and Proposition 3.3 are sharp.

We represent, in Figure 7.1, φ_θ^0 for four different values of θ , with $a = 1$ and $k = 2\pi$. We see that φ_θ^0 is more and more rapidly decaying at infinity as θ increases, in line with (7.2), (3.13), and Proposition 3.3. Then we represent in Figure 7.2 the function

$$(7.3) \quad U_\theta(\varphi_\theta^0)(\mathbf{x}^0) = \int_{\mathbb{R}} h(x_1^0 - a, x_2^0 - \tau_\theta(s)) \varphi_\theta^0(s) \tau_\theta'(s) ds$$

(cf. (4.2)) in the half-space Ω^0 , for $\theta = \pi/6, \pi/4, \pi/3$, evaluating this integral accurately by standard numerical quadrature methods (namely a fifth order composite Gauss quadrature rule on a fine mesh of step length 0.1). From Theorem 3.4 we know that

$$(7.4) \quad U_\theta(\varphi_\theta^0)(\mathbf{x}^0) = u(\mathbf{x}^0) := \frac{i}{4} H_0^{(1)}(k R(x_1^0, x_2^0)), \quad \mathbf{x}^0 \in \Omega_\theta^0,$$

where $\Omega_\theta^0 := \{\mathbf{x}^0 = (x_1^0, x_2^0) : x_1^0 - a > (|x_2^0| - a) \tan \theta\}$. The boundary of Ω_θ^0 is indicated on the figures by dashed lines and, as predicted in (7.4), $U_\theta(\varphi_\theta^0)$ coincides with u in Ω_θ^0 . $U_\theta(\varphi_\theta^0)$ is not equal to u outside Ω_θ^0 ; in particular, it is easy to see from the definition that $U_\theta(\varphi_\theta^0)(a, x_2^0) = 0$ for $|x_2^0| > a$. It appears at first glance that $U_\theta(\varphi_\theta^0)$ is continuous across the dashed lines in Figure 7.2, but an application of the residue theorem, modifying the argument of Theorem 3.4, shows a jump in the value of $U_\theta(\varphi_\theta^0)$ across the dashed lines of

$$\varphi^0(x_2^0 \pm i(x_1^0 - a)) = \varphi_\theta^0(s)$$

at the point $\mathbf{x}^0 = (x_1^0, x_2^0)$ where $x_2^0 = \pm(a + (x_1^0 - a) \cot(\theta)) = \pm a + (s \mp a) \cos(\theta)$. This jump across the dashed lines is just about visible very close to $\mathbf{x}^0 = (a, \pm a)$ but not visible elsewhere because $\varphi_\theta^0(s)$ is exponentially decaying as $s \rightarrow \pm\infty$; see the zoom for $\theta = \pi/3$.

7.2. Discretization of the complex-scaled HSM formulation and validations. We approximate the solution $\Phi_\theta - \Phi_g$ of (3.24) by $\tilde{\Phi}_{\theta, \mathbf{h}} \in \mathbf{V}_{0, \mathbf{h}}$, where $\mathbf{V}_{0, \mathbf{h}}$




FIG. 7.2. Representation of $U_\theta(\varphi_\theta^0)$ in Ω^0 (see (7.3)), where $\varphi_\theta^0(s) = \frac{i}{4} H_0^{(1)}(k R(\tau_\theta(s), a))$, with $a = 1$ and $k = 2\pi$. The dotted lines are part of the boundary of Ω_θ^0 .

is a finite dimensional space $\mathbf{V}_{0,\mathbf{h}} \subset (L_0^2(\mathbb{R}))^4$ that we specify below and $\tilde{\Phi}_{\theta,\mathbf{h}}$ is the solution of the following Galerkin approximation:

$$(7.5) \quad \text{Find } \tilde{\Phi}_{\theta,\mathbf{h}} \in \mathbf{V}_{0,\mathbf{h}} \text{ such that} \\ ((\mathbb{I} - \mathbb{D}_\theta)\tilde{\Phi}_{\theta,\mathbf{h}}, \tilde{\Psi}_{\mathbf{h}})_{(L_0^2(\mathbb{R}))^4} = (\mathbb{D}_\theta\Phi_{g,\mathbf{h}}, \tilde{\Psi}_{\mathbf{h}})_{(L_0^2(\mathbb{R}))^4} \quad \forall \tilde{\Psi}_{\mathbf{h}} \in \mathbf{V}_{0,\mathbf{h}},$$

where \mathbb{D}_θ is defined by (3.25).

To define the approximation space $\mathbf{V}_{0,\mathbf{h}}$, where $\mathbf{h} := (h, q, T)$, let us first introduce $V_{\mathbf{h}} \subset L^2(\mathbb{R})$. To construct $V_{\mathbf{h}}$ we truncate the infinite line at some distance $T > 0$ and build $V_{\mathbf{h}}$ with 1D Lagrange finite elements of degree $q \geq 1$ and maximum element length h supported on $[-T, T]$. The space $\mathbf{V}_{0,\mathbf{h}}$ is nothing else but $\mathbf{V}_{\mathbf{h}} \cap (L_0^2(\mathbb{R}))^4$ where $\mathbf{V}_{\mathbf{h}} := (V_{\mathbf{h}})^4$. Thus, for $\tilde{\Phi}_{\theta,\mathbf{h}} = \{\tilde{\varphi}_{\theta,\mathbf{h}}^0, \tilde{\varphi}_{\theta,\mathbf{h}}^1, \tilde{\varphi}_{\theta,\mathbf{h}}^2, \tilde{\varphi}_{\theta,\mathbf{h}}^3\} \in \mathbf{V}_{0,\mathbf{h}}$, each $\tilde{\varphi}_{\theta,\mathbf{h}}^j$ is a continuous piecewise polynomial function supported in $[-T, T]$ which vanishes on $[-a, a]$. In (7.5), $\Phi_{g,\mathbf{h}} \in \mathbf{V}_{\mathbf{h}} \cap (L^2(-a, a))^4$ is an interpolate of Φ_g . Finally, we approximate Φ_θ by $\Phi_{\theta,\mathbf{h}} = \{\varphi_{\theta,\mathbf{h}}^0, \varphi_{\theta,\mathbf{h}}^1, \varphi_{\theta,\mathbf{h}}^2, \varphi_{\theta,\mathbf{h}}^3\} \in \mathbf{V}_{\mathbf{h}}$, given by $\Phi_{\theta,\mathbf{h}} := \tilde{\Phi}_{\theta,\mathbf{h}} + \Phi_{g,\mathbf{h}}$.

It is clear that the approximation space $\mathbf{V}_{0,\mathbf{h}}$ that we have constructed has the approximation property that, for all $\Phi \in (L_0^2(\mathbb{R}))^4$,

$$\inf_{\tilde{\Psi}_{\mathbf{h}} \in \mathbf{V}_{0,\mathbf{h}}} \|\Phi - \tilde{\Psi}_{\mathbf{h}}\|_{(L_0^2(\mathbb{R}))^4} \rightarrow 0$$

as $h \rightarrow 0$ and $T \rightarrow +\infty$. Thus, and since the sesquilinear form in (7.5) is coercive plus compact on $(L_0^2(\mathbb{R}))^4$ by Theorem 3.7(ii) (cf. Theorem 6.3), standard convergence results for Galerkin methods apply (e.g., [44, Theorem 4.2.9]). These give that, for some $h_0 > 0$ and $T_0 > 0$, the solution $\tilde{\Phi}_{\theta,\mathbf{h}}$ of (7.5) is well-defined for all $0 < h \leq h_0$ and $T \geq T_0$, and a quasi-optimality error estimate holds that, for some constant $C > 0$ and all $0 < h \leq h_0$ and $T \geq T_0$,

$$(7.6) \quad \begin{aligned} \|\Phi_\theta - \Phi_{\theta,\mathbf{h}}\|_{(L^2(\mathbb{R}))^4} &\leq C \inf_{\Psi_{\mathbf{h}} \in \Phi_{g,\mathbf{h}} + \mathbf{V}_{0,\mathbf{h}}} \|\Phi_\theta - \Psi_{\mathbf{h}}\|_{(L^2(\mathbb{R}))^4} \\ &\leq C \left(\|\Phi_\theta\|_{(L^2(\mathbb{R} \setminus (-T, T)))^4} \right. \\ &\quad \left. + \inf_{\Psi_{\mathbf{h}} \in \Phi_{g,\mathbf{h}} + \mathbf{V}_{0,\mathbf{h}}} \|\Phi_\theta - \Psi_{\mathbf{h}}\|_{(L^2(-T, T))^4} \right). \end{aligned}$$

This right-hand side tends to zero as $h \rightarrow 0$ and $T \rightarrow +\infty$, i.e., our Galerkin method is convergent, as long as $\|\Phi_g - \Phi_{g,\mathbf{h}}\|_{(L^2(-a, a))^4} \rightarrow 0$ as $h \rightarrow 0$.

To implement the Galerkin method (7.5), the integrals

$$\int_a^T \left(\int_{-T}^T h(\tau_\theta(t) - a, a - \tau_\theta(s)) \varphi_{\theta, \mathbf{h}}^j(s) \tau_\theta'(s) ds \right) \tilde{\psi}_{\mathbf{h}}^{j\pm 1}(t) dt, \quad \tilde{\psi}_{\mathbf{h}}^{j\pm 1} \in V_{\mathbf{h}},$$

which appear in the variational formulation, need to be approximated. In the results below we use a standard quadrature formula, without any specific treatment of the singularity.

To validate the method, we consider u given by (7.1) which is the solution of (3.24)–(1.2) with $g := u|_{\Sigma_a}$. In the case that $\theta = \pi/6$, $a = 1$, and $k = 2\pi$, we draw in the right-hand side of Figure 7.1 the exact complex-scaled trace φ_θ^0 of u (by symmetry, the four traces are equal in this case) and the computed complex-scaled trace $\varphi_{\theta, \mathbf{h}}^0$, obtained by solving (7.5) with $h = 0.1$, $q = 1$, and $T = 5$. We observe a very good agreement.

More quantitatively, to explore the dependence of the error on T , we plot in Figure 7.3, for a fixed small value of h , the error

$$(7.7) \quad \|\varphi_\theta^0 - \varphi_{\theta, \mathbf{h}}^0\|_{L^2(\mathbb{R})}$$

as a function of T for $\theta = \pi/6, \pi/4, \pi/3$. As $\varphi_{\theta, \mathbf{h}}^0$ is zero outside $(-T, T)$ the error cannot be smaller than the L^2 norm of φ_θ^0 on $\mathbb{R} \setminus (-T, T)$, which decreases like $e^{-k \sin(\theta)T}/T^{1/2}$ as T tends to $+\infty$ by (7.2) and Proposition 3.3. On the other hand, when h is small enough so that the second term on the right-hand side of (7.6) is negligible, the quasi-optimality bound (7.6) and Proposition 3.3 imply that $e^{-k \sin(\theta)T}/T^{1/2}$ is also an upper bound for the error, precisely that the error (7.7) is $\leq C e^{-k \sin(\theta)T}/T^{1/2}$, for some constant $C > 0$ and all sufficiently large T . And, indeed, we observe in Figure 7.3 this rate of exponential behavior as T increases, until the other sources of error become significant.

Using a standard numerical quadrature applied to formula (3.15), with φ_θ^j approximated by $\varphi_{\theta, \mathbf{h}}^0$, we represent finally in Figure 7.4 the numerical solution in $\Omega_\theta^0 \cup \Omega_\theta^1$ (left) and the whole of Ω (right) for $\theta = \pi/6$. To reconstruct the solution in the whole of Ω several choices are possible, since $\theta < \pi/4$ is such that the reconstruction




FIG. 7.3. Absolute error in $\varphi_{\theta, \mathbf{h}}^0$ for three different values of θ computed with P_3 elements ($q = 3$), $a = 1$, and $h = 0.002$ and $k = \pi$ (left), $h = 0.001$, and $k = 2\pi$ (right).

domains overlap (Ω_θ^j overlaps with $\Omega_\theta^{j\pm 1}$, for $j \in \llbracket 0, 3 \rrbracket$). Here we have reconstructed the solution using the identity

$$u(\mathbf{x}^j) = U_\theta(\varphi_\theta^j)(\mathbf{x}^j), \quad \mathbf{x}^j \in \Omega_{\pi/4}^j, \quad j \in \llbracket 0, 3 \rrbracket,$$

where U_θ is defined in (4.2), and where we have used that $\Omega_{\pi/4}^j \subset \Omega_\theta^j$. We notice in Figure 7.4 that these different representations are compatible, up to a small discretization error not visible in the plots.




FIG. 7.4. Reconstruction of the computed solution in $\Omega_\theta^0 \cup \Omega_\theta^1$ (left) and in the whole domain (right), with $\theta = \pi/6$, $a = 1$, $k = 2\pi$.

Finally, we validate formula (4.10) for the far-field pattern. In the present case, the far-field pattern is the same in all directions and one has for $j \in \llbracket 0, 3 \rrbracket$

$$(7.8) \quad F(\cos(j\pi/2), \sin(j\pi/2)) = \frac{1-i}{\sqrt{k\pi}} = \sqrt{\frac{k}{\pi}} \frac{1-i}{2} \int_{\mathbb{R}} \varphi_\theta^j(s) \tau'_\theta(s) \, ds.$$

In Figure 7.5 we plot the real and the imaginary part of the right-hand side of (7.8) when $j = 0$, $a = 1$, and $k = 2\pi$, with φ_θ^0 approximated by $\varphi_{\theta,h}^0$, that is, we plot

$$(7.9) \quad F_{\theta,h} := \sqrt{\frac{k}{\pi}} \frac{1-i}{2} \int_{-T}^T \varphi_{\theta,h}^j(s) \tau'_\theta(s) \, ds$$

as a function of T for different values of θ . We again observe a rapid convergence toward the exact value as T increases.

7.3. Numerical results for the general case. Finally, to discretize problem (6.7), we combine the previous tools with a classical Lagrange finite element approximation of the 2D unknown u_b . Precisely, we use the HSM method to solve the problem of diffraction of the incident plane wave

$$u^i(\mathbf{x}) = \exp(ik(x_1 \cos(\pi/6) + x_2 \sin(\pi/6)))$$

by a perfectly reflecting scatterer which is the union of a disk and a triangle, this union contained in the square Ω_a with $a = 0.8$. The scattering problem is (1.1)–(1.2)




FIG. 7.5. Real (left figure) and imaginary (right figure) parts of the far-field coefficient for three different θ with P3 elements ($q = 3$), $h = 0.001$, $a = 1$, and $k = 2\pi$, computed using the formula (7.9). The black lines indicate the exact values.

in Ω , the domain exterior to the scatterer, with $f = 0$, $\rho = 1$, and the Dirichlet boundary condition $u = g := -u^i$ on $\partial\Omega$. The HSM problem we solve is a variation on (6.7) in which (i) we replace Ω_b by $\tilde{\Omega}_b := \Omega \cap \Omega_b$; (ii) we replace $H^1(\Omega_b)$ by an affine subspace of $H^1(\tilde{\Omega}_b)$ that respects the Dirichlet boundary condition; precisely, we seek $u_b \in \{v \in H^1(\tilde{\Omega}_b) : v = g \text{ on } \partial\Omega\}$. It is easy to see, by a straightforward variation of the arguments of Proposition 6.1 and Theorem 6.3, that this modification of (6.7) remains well-posed: uniqueness holds and the modified sesquilinear form is coercive plus compact on $(L_0^2(\mathbb{R}))^4 \times H_D^1(\tilde{\Omega}_b)$, where $H_D^1(\tilde{\Omega}_b) := \{v \in H^1(\tilde{\Omega}_b) : v = 0 \text{ on } \partial\Omega\}$. As in section 7.2, this implies convergence of Galerkin methods of numerical solution, provided the sequence of approximation spaces used is asymptotically dense in $(L_0^2(\mathbb{R}))^4 \times H_D^1(\tilde{\Omega}_b)$.

In our numerical implementation of a Galerkin method for this HSM formulation we fix $a = 0.8$ and $b = 1.2$, we use P2 elements for both the 2D unknown u_b and the 1D unknowns φ_θ^j , $j \in \llbracket 0, 3 \rrbracket$, with a maximum element diameter for both meshes of $h = 0.05$, choose the truncation parameter $T = 5$, and choose $\theta = \pi/6$. The additional integrals, involving the operator Λ_θ , which appear in the variational formulation are approximated, like the other ones, by a standard quadrature formula.

To reconstruct the solution everywhere several choices are possible, since $\theta < \pi/4$ and the different reconstruction domains overlap (for instance, $\tilde{\Omega}^b$ with Ω_θ^0 , or Ω_θ^0 with Ω_θ^1). We have reconstructed the solution as

$$\begin{aligned} u(\mathbf{x}) &= u_b(\mathbf{x}), \quad \mathbf{x} \in \tilde{\Omega}_b, \\ u(\mathbf{x}^j) &= U_\theta(\varphi_\theta^j)(\mathbf{x}^j), \quad \mathbf{x}^j \in \Omega_{\pi/4}^j \setminus \tilde{\Omega}_b, \quad j \in \llbracket 0, 3 \rrbracket, \end{aligned}$$

where U_θ is defined in (4.2) and where we have used that $\Omega_{\pi/4}^j \subset \Omega_{\pi/6}^j$. See Figure 7.6 for the reconstruction in $\tilde{\Omega}_b \cup \Omega_{\pi/4}^0 \cup \Omega_{\pi/4}^1$ (left), the reconstruction in the whole domain Ω (middle), and the corresponding total field $u + u^i$ (right). We notice that the different deformed half-space representations (the representations in $\Omega_{\pi/4}^j \setminus \tilde{\Omega}_b$, $j \in \llbracket 0, 3 \rrbracket$) are compatible between themselves and are also compatible with the 2D solution u_b in $\tilde{\Omega}_b$.




FIG. 7.6. The diffracted field u reconstructed in $\Omega^b \cup \Omega_{\pi/4}^0 \cup \Omega_{\pi/4}^1$ (left) and in the whole of Ω (middle), and the corresponding total field $u + u^i$ (right) for the case of scattering by a Dirichlet obstacle.

8. Perspectives. Our new complex-scaled HSM method has been presented in this paper for a relatively simple configuration. We expect that it can be extended easily to more complex problems for which the relevant half-space Green's functions are known sufficiently explicitly, such as acoustic scattering in stratified media, including cases where the stratification is different in different half-spaces (see, e.g., [42] for a presentation of the method in the dissipative case). The method is also expected to work well in at least some cases with infinite boundaries, for instance, scattering by an infinite wedge with Dirichlet, Neumann, or Robin boundary conditions. In all these cases, the complex-scaled HSM should be a convenient way to take into account possible surface/guided waves that propagate toward infinity. Elastic scattering in isotropic media can also be considered. More challenging extensions are to the cases where PMLs are observed to fail, such as anisotropic media. A potential advantage over PML of the complex-scaled HSM method in such cases is that it requires the existence of exponentially decaying analytical continuation of the traces of the solution only in a few directions (on the boundaries of a few half-planes).

Appendix A. Properties of the operators \mathbb{D} and \mathbb{D}_θ . The operator \mathbb{D} defined by (2.32) can be rewritten as

$$\mathbb{D} = DS \otimes \mathbb{J} + SD \otimes \mathbb{J}^* \quad \text{with} \quad \mathbb{J} := \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix},$$

where the operator $S \in \mathcal{L}(L^2(\mathbb{R}))$ (the space of continuous linear operators on $L^2(\mathbb{R})$) is defined in (2.20) and the operator $D \in \mathcal{L}(L^2(\mathbb{R}))$ is defined in (2.21) and (2.27). Here $A \otimes \mathbb{M}$ denotes the tensor product of an operator $A \in \mathcal{L}(L^2(\mathbb{R}))$ with a 4×4 scalar matrix \mathbb{M} (see, e.g., [5, section 12.4]), which yields an operator of $\mathcal{L}((L^2(\mathbb{R}))^4)$. But it is actually enough to see this as a simple notation which makes the writing of the proof below easier: $A \otimes \mathbb{M}$ is the block operator matrix obtained by multiplying each scalar component of \mathbb{M} by the operator A . One can easily verify that it satisfies the basic property $\|A \otimes \mathbb{M}\| \leq \|A\| \|\mathbb{M}\|$.

The operator \mathbb{D}_θ , given by (3.25), has the same definition as \mathbb{D} just replacing D by D_θ . The operators D and D_θ satisfy similar properties, given respectively in Proposition 2.1 and in Propositions 3.8, 3.9, and 3.10.

We show in this appendix properties (i) and (ii) of Theorems 2.2 and 3.7. These results are properties of \mathbb{D} and \mathbb{D}_θ , respectively, that are based on the above properties of D and D_θ . Let us give now the proof for (i) and (ii) of Theorem 2.2 for \mathbb{D} ; the same proof holds for Theorem 3.7 for \mathbb{D}_θ .

As D and S are continuous operators on $L^2(\mathbb{R})$, the continuity of \mathbb{D} on $(L^2(\mathbb{R}))^4$ (part (i) of Theorem 2.2) is obvious.

To show part (ii) of Theorem 2.2 we consider \mathbb{D} as an operator on $(L_0^2(\mathbb{R}))^4$. Let us denote by χ_+ (respectively, χ_-) the characteristic function of $(a, +\infty)$ (respectively, $(-\infty, -a)$). We have, by using (2.26), that for $\varphi \in L_0^2(\mathbb{R})$

$$\varphi = \chi_+ \varphi + \chi_- \varphi,$$

and

$$(A.1) \quad \|\varphi\|_{L^2(\mathbb{R})}^2 = \|\chi_+ \varphi\|_{L^2(\mathbb{R})}^2 + \|\chi_- \varphi\|_{L^2(\mathbb{R})}^2.$$

We can reformulate items (ii) and (iii) of Proposition 2.1 using these characteristic functions as follows:

- (ii) $D\chi_+ = L\chi_+ + K\chi_+ \in \mathcal{L}(L^2(\mathbb{R}), L^2(a, +\infty))$, where $L, K \in \mathcal{L}(L^2(a, +\infty))$ are such that $\|L\| \leq 1/\sqrt{2}$ and K is compact.
- (iii) $D\chi_- \in \mathcal{L}(L^2(\mathbb{R}), L^2(a, +\infty))$ is compact.

These properties, and that $S\chi_\pm = \chi_\mp S$, lead us to decompose \mathbb{D} as an operator on $(L_0^2(\mathbb{R}))^4$ as

$$\mathbb{D} = \mathbb{L} + \mathbb{K},$$

where

$$\mathbb{K} := (DS\chi_+ + KS\chi_-) \otimes \mathbb{J} + (SD\chi_- + SK\chi_+) \otimes \mathbb{J}^*$$

and

$$\mathbb{L} := (LS\chi_-) \otimes \mathbb{J} + (SL\chi_+) \otimes \mathbb{J}^*.$$

It follows from (ii) and (iii) that \mathbb{K} is compact. Moreover, noting that $\chi^+ L = L$, so that $\chi^- S L = S L$, we have

$$\mathbb{L} = \chi_+ (LS \otimes \mathbb{J}) \chi_- + \chi_- (SL \otimes \mathbb{J}^*) \chi_+.$$

We deduce then by (A.1) that, for all $\Phi \in (L_0^2(\mathbb{R}))^4$,

$$\begin{aligned} \|\mathbb{L}\Phi\|_{L^2(\mathbb{R})^4}^2 &= \|(LS \otimes \mathbb{J}) \chi_- \Phi\|_{L^2(\mathbb{R})^4}^2 + \|(SL \otimes \mathbb{J}^*) \chi_+ \Phi\|_{L^2(\mathbb{R})^4}^2 \\ &\leq \frac{1}{2} \|\chi_- \Phi\|_{L^2(\mathbb{R})^4}^2 + \frac{1}{2} \|\chi_+ \Phi\|_{L^2(\mathbb{R})^4}^2 = \frac{1}{2} \|\Phi\|_{L^2(\mathbb{R})^4}^2, \end{aligned}$$

where we have used that $\|LS \otimes \mathbb{J}\| \leq \|LS\| \|\mathbb{J}\| \leq 1/\sqrt{2}$, and the same bound for $SL \otimes \mathbb{J}^*$.

Appendix B. Technical lemmas. The lemmas in this annex (cf. [22, Lemma 4.4]) concern the complex functions

$$(B.1) \quad R(\hat{z}, z) := (\hat{z}^2 + z^2)^{1/2}, \quad \hat{R}(\hat{z}, z) := (\hat{z} + z^2)^{1/2}, \quad z, \hat{z} \in \mathbb{C}.$$

Note that, as throughout the rest of the paper, all square roots in this appendix are principal square roots, i.e., square roots with argument in the range $(-\pi/2, \pi, 2]$.

LEMMA B.1. *We have*

$$|R(t, z)|^2 \geq |\cos(\operatorname{Arg}(z))| (t^2 + |z|^2), \quad t \in \mathbb{R}, z \in \mathbb{C}, z \neq 0.$$

Proof. Let $\gamma := \operatorname{Arg}(z)$. We have

$$|R(t, z)|^4 = |t^2 + |z|^2 e^{2i\gamma}|^2 = t^4 + |z|^4 + 2t^2|z|^2 \cos(2\gamma) = (t^2 - |z|^2)^2 + 4t^2|z|^2 \cos^2(\gamma),$$

which yields

$$|R(t, z)|^4 \geq \cos^2(\gamma) \left[(t^2 - |z|^2)^2 + 4t^2|z|^2 \right] = \cos^2(\gamma) (t^2 + |z|^2)^2. \quad \square$$

LEMMA B.2. *There exists a constant $C > 0$ such that, for all $z \in \mathbb{C}$ and $t \in \mathbb{R}$,*

$$|R(t, z) - z| \leq C \frac{t^2}{|z|}, \quad \Re(z) > 0, |z| \geq |t|;$$

in particular

$$\Im(R(t, z)) \geq \Im(z) - C \frac{t^2}{|z|}, \quad \Re(z) > 0, |z| \geq |t|.$$

Moreover, for all $A > 0$, there exists a constant $C' > 0$ such that, for $-A \leq t \leq A$,

$$\Im(R(t, z)) \geq \Im(z) - C', \quad \Re(z) > 0.$$

Proof. Since the function $z \mapsto (1 + z)^{1/2} - 1$ is an analytic function of z in the open unit disk that vanishes at the origin and is bounded in the closed disk, for some constant $C > 0$,

$$(B.2) \quad |(1 + z)^{1/2} - 1| \leq C|z|, \quad |z| \leq 1.$$

Further, the function

$$z \mapsto (t^2 + z^2)^{1/2} - z \left(1 + \frac{t^2}{z^2} \right)^{1/2}$$

is analytic in the domain $\{z \in \mathbb{C} : \Re(z) > 0\}$ and it vanishes for $z \in \mathbb{R}$ with $z > 0$, so that it vanishes everywhere in $\{z \in \mathbb{C} : \Re(z) > 0\}$. This implies that

$$R(t, z) - z = z \left[\left(1 + \frac{t^2}{z^2} \right)^{1/2} - 1 \right], \quad \Re(z) > 0.$$

Combined with (B.2), this proves the first inequality.

The second inequality is a direct consequence of the first since

$$|R(t, z) - z| \geq \Im(z - R(t, z)).$$

Finally, the second inequality yields that, for any constant $A > 0$ and for all $|t| \leq A \leq |z|$, $\Im(R(t, z)) \geq \Im(z) - CA$. But a similar inequality also holds for $|z| \leq A$, possibly with another constant, since $|R(t, z)| \leq \sqrt{2}A$ for $|t| \leq A$ and $|z| \leq A$. \square

Remark B.3. Note that, replacing z by $-z$, one directly deduces from the previous lemma similar results valid for $\Re(z) < 0$. In particular,

$$|R(t, z) + z| \leq C \frac{t^2}{|z|}, \quad \Re(z) < 0, |z| \geq |t|,$$

and

$$\Im(R(t, z)) \geq -\Im(z) - C \frac{t^2}{|z|}, \quad \Re(z) < 0, |z| \geq |t|.$$

Analogous results can be proved in the more general case where the positive real number t^2 is replaced by any complex number \hat{z} .

LEMMA B.4. *There exists a constant $C > 0$ such that, for all $z, \hat{z} \in \mathbb{C}$, $z \neq 0$,*

$$(B.3) \quad \begin{aligned} |\hat{R}(\hat{z}, z) - z| &\leq C \left| \frac{\hat{z}}{z} \right|, \quad \Re(z) \geq |\hat{z}|^{1/2}, \\ \Im(\hat{R}(\hat{z}, z)) &\geq \Im(z) - C \left| \frac{\hat{z}}{z} \right|, \quad \Re(z) \geq |\hat{z}|^{1/2}. \end{aligned}$$

Moreover, if U is a bounded subset of \mathbb{C} and $0 < \gamma < \pi/2$, there exists a constant $C' > 0$ such that

$$\Im(\hat{R}(\hat{z}, z)) \geq \Im(z) - C', \quad \hat{z} \in U, |\arg(z)| < \gamma.$$

Proof. To proceed as in the proof of Lemma B.2, we just have to show that the function

$$z \mapsto (\hat{z} + z^2)^{1/2} - z \left(1 + \frac{\hat{z}}{z^2} \right)^{1/2}$$

is analytic in the domain $\{\Re(z) > |\hat{z}|^{1/2}\}$. For that, we have to check that the branch cuts of the functions $z \mapsto (\hat{z} + z^2)^{1/2}$ and $z \mapsto (1 + \hat{z}/z^2)^{1/2}$ do not intersect the domain $\Re(z) > |\hat{z}|^{1/2}$.

This is clear for the function $z \mapsto (1 + \hat{z}/z^2)^{1/2}$. The branch cut of the function $z \mapsto (\hat{z} + z^2)^{1/2}$ is the subset of the hyperbola $\{z \in \mathbb{C} : \Im(z^2 + \hat{z}) = 0\}$ where $\Re(z^2 + \hat{z}) \leq 0$. The intersection of this hyperbola with the domain $\Re(z) > |\hat{z}|^{1/2}$ is the connected set

$$\hat{H} = \{z \in \mathbb{C} : \Im(z^2 + \hat{z}) = 0, \Re(z) > |\hat{z}|^{1/2}\},$$

which describes a curve which is asymptotic to the real axis when $\Re(z) \rightarrow +\infty$. To conclude, we have to prove that for all $z \in \hat{H}$, $\Re(z^2 + \hat{z}) > 0$. This is clearly true for large values of $\Re(z)$. If it were not true for all $z \in \hat{H}$, there would exist some $z_0 \in \hat{H}$ such that $\Re(z_0^2 + \hat{z}) = 0$. But then $z_0^2 + \hat{z} = 0$, so that $|z_0| = |\hat{z}|^{1/2}$, which is impossible since $\Re(z_0) > |\hat{z}|^{1/2}$. \square

An important application of the previous lemmas for our purpose is the following.

LEMMA B.5. *For $\theta \in (0, \pi/2)$, let τ_θ be defined by (3.11), and suppose that $\theta_0 \in [\theta, \pi/2)$. Then, for $w \in \mathbb{C}$ such that $w \neq 0$ and $-\theta_0 + \theta \leq \arg(w) \leq \theta_0$, and for all $s \in \mathbb{R}$, it holds that*

$$(B.4) \quad |R(w, \tau_\theta(s) - a)|^2 \geq \cos(\theta_0)(|w|^2 + |\tau_\theta(s) - a|^2) \geq \cos^2(\theta_0)(|w|^2 + |\tau_\theta(s) - a|^2).$$

Further, there exists a constant $C > 0$, depending only on a and θ , such that, for all $s \in \mathbb{R}$,

$$(B.5) \quad \Im(R(w, \tau_\theta(s) - a)) \geq [\cos(\theta - \text{Arg}(w))]^{1/2} \Im(w) - C \geq \cos(\theta - \text{Arg}(w)) \Im(w) - C,$$

provided $w \neq 0$ and $0 \leq \text{Arg}(w) \leq \theta$.

Proof. Let $\gamma := \text{Arg}(w) \in [\theta - \theta_0, \theta_0]$. We have, for all $s \in \mathbb{R}$,

$$|R(w, \tau_\theta(s) - a)| = |R(|w|e^{i\gamma}, \tau_\theta(s) - a)| = |R(|w|, (\tau_\theta(s) - a)e^{-i\gamma})|.$$

Applying Lemma B.1 we obtain the inequality

$$(B.6) \quad |R(w, \tau_\theta(s) - a)|^2 \geq |\cos(\hat{\gamma}(s))|(|w|^2 + |\tau_\theta(s) - a|^2),$$

where $\hat{\gamma}(s) := \text{Arg}((\tau_\theta(s) - a)e^{-i\gamma})$.

We now consider separately the three cases $|s| \leq a$, $s > a$, and $s < -a$ to derive the two inequalities of the lemma.

Case 1. For $|s| \leq a$, since $\tau_\theta(s) = s$, we have $|\cos(\hat{\gamma}(s))| = \cos(\gamma) \geq \cos(\theta_0)$ so that (B.4) holds. Further, applying the third inequality of Lemma B.2 we get that there exists a constant $c > 0$, dependent only on a , such that

$$(B.7) \quad \Im(R(w, \tau_\theta(s) - a)) = \Im(R(s - a, w)) \geq \Im(w) - c.$$

Case 2. For $s > a$, since $\tau_\theta(s) - a = \tilde{s}e^{i\theta}$, where $\tilde{s} := s - a > 0$, and $\hat{\gamma}(s) = \theta - \gamma$ so that $|\cos(\hat{\gamma}(s))| = \cos(\theta - \gamma) \geq \cos(\theta_0)$, we have from (B.6) that

$$(B.8) \quad |R(w, \tau_\theta(s) - a)|^2 = |R(w, \tilde{s}e^{i\theta})|^2 \geq \cos(\theta - \gamma)(|w|^2 + \tilde{s}^2),$$

so that (B.4) holds. To see that (B.5) holds, for $\gamma \in [0, \theta]$ as required in the lemma, note that

$$(B.9) \quad \text{Arg}(R(w, \tau_\theta(s) - a)) = \text{Arg}(R(|w|e^{i\gamma}, \tilde{s}e^{i\theta})) \in [\gamma, \theta].$$

Combining this with the bound (B.8) we see that

$$(B.10) \quad \begin{aligned} \Im(R(w, \tau_\theta(s) - a)) &= \Im(R(w, \tilde{s}e^{i\theta})) = |R(w, \tilde{s}e^{i\theta})| \sin(\text{Arg}(R(w, \tilde{s}e^{i\theta}))) \\ &\geq \sqrt{\cos(\theta - \gamma)} |w| \sin(\gamma) = \sqrt{\cos(\theta - \gamma)} \Im(w). \end{aligned}$$

Case 3. For $s < -a$, setting $\tilde{s} := -(s + a) > 0$, we have $\tau_\theta(s) - a = -(\tilde{s}e^{i\theta} + 2a)$ and $|\cos(\hat{\gamma}(s))| = |\cos(\hat{\theta}(\tilde{s}) - \gamma)|$, where $\hat{\theta}(\tilde{s}) := \text{Arg}(\tilde{s}e^{i\theta} + 2a)$. Since $0 \leq \hat{\theta}(\tilde{s}) \leq \theta$ for $\tilde{s} > 0$, and $\gamma \in [\theta - \theta_0, \theta_0]$, we have

$$|\cos(\hat{\gamma}(s))| \geq \min_{0 \leq \tilde{\gamma} \leq \theta} |\cos(\gamma - \tilde{\gamma})| \geq \cos(\theta_0),$$

so that (B.4) follows from (B.6).

To show (B.5), for $0 \leq \gamma \leq \theta$ as required, we rewrite

$$(B.11) \quad \begin{aligned} R(w, \tilde{s}e^{i\theta} + 2a) &= [R(w, \tilde{s}e^{i\theta})^2 + 4a^2 + 4a\tilde{s}e^{i\theta}]^{1/2} \\ &= \hat{R}(4a^2 + 4a\tilde{s}e^{i\theta}, R(w, \tilde{s}e^{i\theta})), \end{aligned}$$

where \hat{R} is defined by (B.1). Using (B.8) and (B.9) we see that

$$|R(w, \tilde{s}e^{i\theta})|^2 \geq \cos(\theta - \gamma)\tilde{s}^2 \geq \cos(\theta)\tilde{s}^2 \quad \text{and} \quad \text{Arg}(R(w, \tilde{s}e^{i\theta})) \in [\gamma, \theta].$$

Thus, by (B.3) applied to (B.11), there exist constants $C > 0$ and $\tilde{s}_0 > 0$, depending only on θ and a , such that

$$\Im(R(w, \tilde{s}e^{i\theta} + 2a)) \geq \Im(R(w, \tilde{s}e^{i\theta})) - C \frac{1 + \tilde{s}}{\tilde{s}},$$

for $\tilde{s} \geq \tilde{s}_0$. Hence, and using (B.10), it follows that, for $\tilde{s} \geq \max(1, \tilde{s}_0)$,

$$\Im(R(w, \tilde{s}e^{i\theta} + 2a)) \geq \sqrt{\cos(\theta - \gamma)} \Im(w) - 2C.$$

On the other hand, for $\tilde{s} \leq \max(1, \tilde{s}_0)$ the last inequality of Lemma B.4 gives that there exists a constant $C' > 0$, depending only on a , such that

$$\Im(R(w, \tilde{s}e^{i\theta} + 2a)) \geq \Im(w) - C'.$$

Gathering the two estimates we have that, for $s < -a$,

$$(B.12) \quad \Im(R(w, \tau_\theta(s) - a)) \geq \sqrt{\cos(\theta - \gamma)} \Im(w) - \max(C', 2C).$$

We have shown that (B.4) holds in each case. That (B.5) also holds follows from (B.7), (B.10), and (B.12), trivially noting that $(\cos(t))^{1/2} \geq \cos(t)$ for $t \in \mathbb{R}$. \square

Appendix C. Mapping properties of complex-scaled integral operators.

In this appendix we prove mapping properties of the analytic continuations into the complex plane of the single- and double-layer potential operators \mathcal{S}^j and \mathcal{D}^j , defined by (3.5) for $j \in \llbracket 0, 3 \rrbracket$, $\phi \in L^2(\Sigma_a)$, and $|t| > a$. The proofs use the bounds established in Appendix B and bounds on the relevant Hankel functions. Specifically [35, Lemma 3.4], for some constant $c_1 > 0$,

$$(C.1) \quad \left| e^{-iz} H_1^{(1)}(z) \right| \leq c_1 \left(|z|^{-1} + (1 + |z|)^{-1/2} \right), \quad \Re(z) > 0.$$

Similarly, it follows from (3.6) and [30, section 10.2(ii)] that, for some constant $c_0 > 0$,

$$(C.2) \quad \left| e^{-iz} H_0^{(1)}(z) \right| \leq c_0 M(|z|), \quad \Re(z) > 0,$$

where $M(t) := \log(1 + t^{-1}) + (1 + t)^{-1/2}$, for $t > 0$.

PROPOSITION C.1. *For every $\theta \in (0, \pi/2)$ there exists some constant $C > 0$, that depends only on a , k , and θ , such that, for $j \in \llbracket 0, 3 \rrbracket$ and every $\phi \in L^2(\Sigma_a)$,*

$$(C.3) \quad |\mathcal{S}^j \phi(z)| \leq C M(|z - a|) \exp(-k \Im(z)) \|\phi\|_{L^2(\Sigma_a)},$$

$$(C.4) \quad |\mathcal{D}^j \phi(z)| \leq C |z - a|^{-1/2} \exp(-k \Im(z)) \|\phi\|_{L^2(\Sigma_a)},$$

for $\Re(z) > a$ with $|\arg(z - a)| \leq \theta$, while

$$(C.5) \quad |\mathcal{S}^j \phi(z)| \leq C M(|z + a|) \exp(k \Im(z)) \|\phi\|_{L^2(\Sigma_a)},$$

$$(C.6) \quad |\mathcal{D}^j \phi(z)| \leq C |z + a|^{-1/2} \exp(k \Im(z)) \|\phi\|_{L^2(\Sigma_a)},$$

for $\Re(z) < -a$ with $|\arg(-z - a)| \leq \theta$.

Proof. We prove only the bounds (C.3)–(C.4); the proofs of (C.5)–(C.6) are identical. Throughout this proof C will denote any positive constant depending only on a , k , and θ , not necessarily the same at each occurrence, and we assume that $\mathbf{y}^j \in \Sigma_a$ and that $\Re(z) > a$ with $|\arg(z - a)| \leq \theta$.

It follows from (the analytic continuation of) (3.5) and the Cauchy–Schwarz inequality that

$$|\mathcal{S}^j \phi(z)| \leq C I_S^{1/2} \|\phi\|_{L^2(\Sigma_a)}, \quad |\mathcal{D}^j \phi(z)| \leq C I_D^{1/2} \|\phi\|_{L^2(\Sigma_a)},$$

where

$$I_S := \int_{\Sigma_a} |\Phi(\mathbf{x}^j(z), \mathbf{y}^j)|^2 ds(\mathbf{y}^j), \quad I_D := \int_{\Sigma_a} \left| \frac{\partial \Phi(\mathbf{x}^j(z), \mathbf{y}^j)}{\partial n(\mathbf{y}^j)} \right|^2 ds(\mathbf{y}^j).$$

Now, from (C.2), and since $\mathbf{x}^j(z) := (a, z)$ and recalling the definition (2.15),

$$|\Phi(\mathbf{x}^j(z), \mathbf{y}^j)| \leq C M \left(k |R(a - y_1^j, z - y_2^j)| \right) \left| \exp(ikR(a - y_1^j, z - y_2^j)) \right|.$$

Further, it follows from Lemma B.2 that, for some constant $C_a > 0$ depending only on a ,

$$(C.7) \quad |\exp(ikR(a - y_1^j, z - y_2^j))| \leq \exp(-k(\Im(z) - C_a)) \leq C \exp(-k\Im(z))$$

and from Lemma B.1 that

$$(C.8) \quad |R(a - y_1^j, z - y_2^j)| \geq \cos(\theta)((a - y_1^j)^2 + |z - a|^2)^{1/2} \geq \cos(\theta)|z - a|.$$

Thus, noting that $M(t)$ is decreasing as t increases and that, for every $c > 0$, $t \mapsto M(ct)/M(t)$ is a bounded function on $t > 0$, it follows that

$$|\Phi(\mathbf{x}^j(z), \mathbf{y}^j)| \leq CM(|z - a|) \exp(-k\Im(z)),$$

so that $I_S^{1/2} \leq CM(|z - a|) \exp(-k\Im(z))$ and (C.3) follows.

Writing $n(\mathbf{y}^j) = (n_1(\mathbf{y}^j), n_2(\mathbf{y}^j))$ in the (x_1^j, x_2^j) coordinate system, we see that

$$(C.9) \quad \begin{aligned} & \left| \frac{\partial \Phi(\mathbf{x}^j(z), \mathbf{y}^j)}{\partial n(\mathbf{y}^j)} \right| \\ &= \frac{k}{4} \left| H_1^{(1)}(kR(a - y_1^j, z - y_2^j)) \right| \frac{|n_1(\mathbf{y}^j)(y_1^j - a) + n_2(\mathbf{y}^j)(y_2^j - z)|}{|R(a - y_1^j, z - y_2^j)|}. \end{aligned}$$

How we bound the right-hand side of this equation depends on which side of Σ_a the point \mathbf{y}^j is located. When $\mathbf{y}^j \in \Sigma_a^j \subset \Sigma_a$ the right-hand side of (C.9) vanishes, since then $y_1^j = a$ and $n_2(\mathbf{y}^j) = 0$. When $\mathbf{y}^j \in \partial\Omega \setminus (\Sigma_a^j \cup \Sigma_a^{j+1})$ it holds that $y_1^j = -a$ or $y_2^j = -a$, so that $|R(\mathbf{x}^j(z), \mathbf{y}^j)| \geq C$ by Lemma B.1, from which it follows, applying (C.1) and (C.7), that the right-hand side of (C.9) is bounded by $C \exp(-k\Im(z))$. Thus

$$I_D \leq C \exp(-2k\Im(z)) + I_D^*, \quad \text{where } I_D^* := \int_{\Sigma_a^{j+1}} \left| \frac{\partial \Phi(\mathbf{x}^j(z), \mathbf{y}^j)}{\partial n(\mathbf{y}^j)} \right|^2 ds(\mathbf{y}^j).$$

Finally, when $\mathbf{y}^j \in \Sigma_a^{j+1}$ it holds that $n_1(\mathbf{y}^j) = 0$ and $y_2^j = a$, so that, where $\rho := |z - a|$, (C.9) says that

$$(C.10) \quad \begin{aligned} & \left| \frac{\partial \Phi(\mathbf{x}^j(z), \mathbf{y}^j)}{\partial n(\mathbf{y}^j)} \right| = \frac{k}{4} \left| H_1^{(1)}(kR(a - y_1^j, z - y_2^j)) \right| \frac{\rho}{|R(a - y_1^j, z - y_2^j)|} \\ & \leq C \left(\frac{\rho}{\rho^2 + (a - y_1^j)^2} + (1 + \rho)^{-1/2} \right) \exp(-k\Im(z)), \end{aligned}$$

using (C.1), (C.7), and (C.8). Thus

$$e^{2k\Im(z)} I_{\mathcal{D}}^* \leq \frac{C}{1+\rho} + C \int_{-a}^a \frac{\rho^2 dy_1^j}{(\rho^2 + (y_1^j - a)^2)^2} \leq \frac{C}{\rho},$$

where we see the final bound by substituting $s := (y_1^j - a)/\rho$. Thus $I_{\mathcal{D}}^{1/2} \leq C|z - a|^{-1/2}e^{-k\Im(z)}$ and the bound (C.4) follows. \square

The final result of this appendix, and arguments we make elsewhere in the paper, depend on a mapping property of the classical double-layer potential operator on the boundary of a quadrant. For the convenience of the reader we state this key and well-known result and sketch its proof in the following proposition.

PROPOSITION C.2. *Let h_0 denote the kernel function h defined in (2.14) in the case that $k = 0$, so that $h_0(x_1, x_2) = x_1/(\pi(x_1^2 + x_2^2))$, for $x_1 > 0$ and $x_2 \in \mathbb{R}$. For $\phi \in L^2(0, +\infty)$ define $\mathfrak{D}\phi : (0, +\infty) \rightarrow \mathbb{C}$ by*

$$\mathfrak{D}\phi(s) := \int_0^{+\infty} h_0(s, t) \phi(t) dt = \frac{1}{\pi} \int_0^{+\infty} \frac{s}{s^2 + t^2} \phi(t) dt, \quad s > 0.$$

Then $\mathfrak{D}\phi \in L^2(0, +\infty)$ and the mapping $\mathfrak{D} : L^2(0, +\infty) \rightarrow L^2(0, +\infty)$ is bounded, with norm $\|\mathfrak{D}\| = 1/\sqrt{2}$.

Proof. This result can be proved by making use of the equivalence of (2.22) and (2.23) in the static case $k = 0$ as in [12], or directly via Mellin transform methods (cf. [41, 43]). Equivalently, we observe as in [17] that the mapping $\mathfrak{I} : L^2(0, +\infty) \rightarrow L^2(\mathbb{R})$, given by $\mathfrak{I}\phi(t) = \phi(e^{-t})e^{-t/2}$, $t \in \mathbb{R}$, is unitary, as is the Fourier transform operator $\mathfrak{F} : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$, $\phi \mapsto \widehat{\phi}$, given by (2.9). Further [17], for $\phi \in L^2(\mathbb{R})$,

$$\mathfrak{I}\mathfrak{D}\mathfrak{I}^{-1}\phi(s) = \int_{\mathbb{R}} \kappa(s-t)\phi(t) dt, \quad s \in \mathbb{R},$$

where $\kappa(\tau) := e^{\tau/2}/(\pi(e^{2\tau} + 1))$, for $\tau \in \mathbb{R}$, and [17]

$$\widehat{\kappa}(\xi) = \frac{1}{\sqrt{2\pi}} \frac{\sinh(\pi(\xi - i/2)/2)}{\sinh(\pi(\xi - i/2))}, \quad \xi \in \mathbb{R}.$$

Thus, for $\phi \in L^2(\mathbb{R})$,

$$\mathfrak{F}\mathfrak{I}\mathfrak{D}\mathfrak{I}^{-1}\mathfrak{F}^{-1}\widehat{\phi} = \sqrt{2\pi}\widehat{\kappa}\widehat{\phi},$$

so that [17], since $\widehat{\kappa} \in L^\infty(\mathbb{R})$, \mathfrak{D} is bounded on $L^2(0, +\infty)$ with

$$\|\mathfrak{D}\| = \|\mathfrak{F}\mathfrak{I}\mathfrak{D}\mathfrak{I}^{-1}\mathfrak{F}^{-1}\| = \sqrt{2\pi}\|\widehat{\kappa}\|_{L^\infty(\mathbb{R})} = \sqrt{2\pi}|\widehat{\kappa}(0)| = 1/\sqrt{2}. \quad \square$$

Remark C.3. Let D_0 denote the operator D , given by (2.23) and (2.27), in the case that $k = 0$, so that, for $\phi \in L^2(\mathbb{R})$, $D_0\phi(t) = 0$, for $t \leq a$, while

$$D_0\phi(t) = \int_{\mathbb{R}} h_0(t-a, s-a) \phi(s) ds = \frac{1}{\pi} \int_{\mathbb{R}} \frac{t-a}{(t-a)^2 + (s-a)^2} \phi(t) dt, \quad t > a,$$

where h_0 is as defined in Proposition C.2. Then it is clear from the above proposition that, as an operator on $L^2(a, +\infty)$, D_0 is bounded with norm $\|D_0\| = \|\mathfrak{D}\| = 1/\sqrt{2}$, and that D_0 is also bounded as an operator on $L^2(\mathbb{R})$, with norm $\|D_0\| = 2\|\mathfrak{D}\| = \sqrt{2}$.

PROPOSITION C.4. For $0 < \theta < \pi/2$ and $j \in \llbracket 0, 3 \rrbracket$, \mathcal{S}_θ^j and \mathcal{D}_θ^j are continuous operators from $L^2(\Sigma_a)$ to $L^2(-\infty, -a) \oplus L^2(a, +\infty)$, where, for $\phi \in L^2(\Sigma_a)$,

$$\mathcal{S}_\theta^j \phi(s) := \mathcal{S}^j \phi(\tau_\theta(s)), \quad \mathcal{D}_\theta^j \phi(s) := \mathcal{D}^j \phi(\tau_\theta(s)), \quad |s| > a.$$

Proof. It is clear from the bounds (C.3) and (C.5) that \mathcal{S}_θ^j maps $L^2(\Sigma_a)$ continuously to $L^2(-\infty, -a) \oplus L^2(a, +\infty)$. The analogous bounds (C.4) and (C.6) do not quite imply that $\mathcal{D}_\theta^j \phi \in L^2(-\infty, -a) \oplus L^2(a, +\infty)$ for each $\phi \in L^2(\Sigma_a)$, since $(s-a)^{-1/2} e^{-k \sin(\theta)(s-a)}$ is in $L^1(a, +\infty)$ but not in $L^2(a, +\infty)$. To see that $\mathcal{D}_\theta^j : L^2(\Sigma_a) \rightarrow L^2(a, +\infty)$ and is continuous we argue as in the proof of Proposition C.1, in particular using (C.9) and (C.10), and recalling that, except when $\mathbf{y}^j \in \Sigma_a^{j+1}$, the right-hand side of (C.9) is $\leq C \exp(-k \Im(z))$. These bounds imply that, for $\phi \in L^2(\Sigma_a)$ and $s > a$,

$$|\mathcal{D}_\theta^j \phi(s)| \leq C e^{-k \sin(\theta)(s-a)} \int_{\Sigma_a} |\phi(\mathbf{y}^j)| \, ds(\mathbf{y}^j) + C \int_{\Sigma_a^{j+1}} \frac{|s-a| |\phi(\mathbf{y}^j)|}{(s-a)^2 + (y_1^j - a)^2} \, ds(\mathbf{y}^j),$$

where, throughout the proof, $C > 0$ denotes a constant that depends only on θ , a , and k . Thus

$$|\mathcal{D}_\theta^j \phi(s)| \leq C e^{-k \sin(\theta)(s-a)} \|\phi\|_{L^2(\Sigma_a)} + C D_0 \psi(s),$$

where D_0 is as in Remark C.3 above, while $\psi \in L^2(-a, a)$ denotes the restriction of $|\phi|$ to Σ_a^{j+1} , precisely $\psi(y_1^j) := |\phi((y_1^j, a))|$, for $-a < y_1^j < a$, while $\psi(y_1^j) := 0$, for $|y_1^j| \geq a$. Since (Remark C.3) D_0 is a bounded operator on $L^2(\mathbb{R})$, it follows that

$$\|\mathcal{D}_\theta^j \phi\|_{L^2(a, +\infty)} \leq C \|\phi\|_{L^2(\Sigma_a)} + C \|D_0\| \|\psi\|_{L^2(-a, a)} \leq C \|\phi\|_{L^2(\Sigma_a)}.$$

Arguing in the same way, we see that \mathcal{D}_θ^j is also continuous as a mapping from $L^2(\Sigma_a)$ to $L^2(-\infty, -a)$, and the proof is complete. \square

REFERENCES

- [1] M. ABRAMOWITZ AND I. A. STEGUN, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Dover, New York, 1964.
- [2] J. AGUILAR AND J.-M. COMBES, *A class of analytic perturbations for one-body Schrödinger Hamiltonians*, Comm. Math. Phys., 22 (1971), pp. 269–279.
- [3] H. AMANN AND J. ESCHER, *Analysis III*, Birkhäuser, Basel, 2009.
- [4] T. ARENS AND T. HOHAGE, *On radiation conditions for rough surface scattering problems*, IMA J. Appl. Math., 70 (2005), pp. 839–847.
- [5] J.-P. AUBIN, *Applied Functional Analysis*, 2nd ed., John Wiley & Sons, New York, 2000.
- [6] E. BALSLEV AND J.-M. COMBES, *Spectral properties of many-body Schrödinger operators with dilatation-analytic interactions*, Comm. Math. Phys., 22 (1971), pp. 280–294.
- [7] V. BARONIAN, A.-S. BONNET-BENDHIA, S. FLISS, AND Y. TJANDRAWIDJAJA, *The half-space matching method for Lamb wave scattering in anisotropic plates*, in Proceedings of the 2nd International Conference on Advanced Modelling of Wave Propagation in Solids, Prague, 2018.
- [8] E. BÉCACHÉ, S. FAUQUEUX, AND P. JOLY, *Stability of perfectly matched layers, group velocities and anisotropic waves*, J. Comput. Phys., 188 (2003), pp. 399–433.
- [9] A.-S. BONNET-BEN DHIA, S. N. CHANDLER-WILDE, AND S. FLISS, *On the Half-Space Matching Method for Real Wavenumber*, preprint, arXiv:2111.06793.
- [10] A.-S. BONNET-BEN DHIA, G. DAKHIA, C. HAZARD, AND L. CHORFI, *Diffraction by a defect in an open waveguide: A mathematical analysis based on a modal radiation condition*, SIAM J. Appl. Math., 70 (2009), pp. 677–693.

- [11] A.-S. BONNET-BENDHIA, S. FLISS, AND Y. TJANDRAWIDJAJA, *Numerical analysis of the half-space matching method with Robin traces on a convex polygonal scatterer*, in Maxwell's Equations: Analysis and Numerics, U. Langer, D. Pauly, and S. Repin, eds., De Gruyter, Berlin, 2019, pp. 105–144.
- [12] A.-S. BONNET-BENDHIA, S. FLISS, AND A. TONNOIR, *The halfspace matching method: A new method to solve scattering problem in infinite media*, J. Comput. Appl. Math., 338 (2018), pp. 44–68.
- [13] A.-S. BONNET-BENDHIA, B. GOURSAUD, AND C. HAZARD, *Mathematical analysis of the junction of two acoustic open waveguides*, SIAM J. Appl. Math., 71 (2011), pp. 2048–2071.
- [14] A.-S. BONNET-BENDHIA AND A. TILLEQUIN, *A generalized mode matching method for the junction of open waveguides*, in Proceedings of the 5th International Conference on Mathematical and Numerical Aspects of Wave Propagation, A. Bermúdez, D. Gómez, C. Hazard, P. Joly, and J. E. Roberts, eds., SIAM, Philadelphia, 2000, pp. 399–403.
- [15] A.-S. BONNET-BENDHIA AND A. TILLEQUIN, *A limiting absorption principle for scattering problems with unbounded obstacles*, Math. Methods Appl. Sci., 24 (2001), pp. 1089–1111.
- [16] H. BRAKHAGE AND P. WERNER, *Über das Dirichletsche Aussenraumproblem für die Helmholtzsche Schwingungsgleichung*, Arch. Math., 16 (1965), pp. 325–329.
- [17] G. A. CHANDLER, *Galerkin's method for boundary integral equations on polygonal domains*, ANZIAM J., 26 (1984), pp. 1–13.
- [18] S. CHANDLER-WILDE AND B. ZHANG, *Electromagnetic scattering by an inhomogeneous conducting or dielectric layer on a perfectly conducting plate*, Proc. A, 454 (1998), pp. 519–542.
- [19] S. N. CHANDLER-WILDE, *The impedance boundary value problem for the Helmholtz equation in a half-plane*, Math. Methods Appl. Sci., 20 (1997), pp. 813–840.
- [20] S. N. CHANDLER-WILDE AND J. ELSCHNER, *Variational approach in weighted Sobolev spaces to scattering by unbounded rough surfaces*, SIAM J. Math. Anal., 42 (2010), pp. 2554–2580.
- [21] S. N. CHANDLER-WILDE, I. G. GRAHAM, S. LANGDON, AND E. A. SPENCE, *Numerical-asymptotic boundary integral methods in high-frequency acoustic scattering*, Acta Numer., 21 (2012), pp. 89–305.
- [22] S. N. CHANDLER-WILDE, D. P. HEWETT, S. LANGDON, AND A. TWIGGER, *A high frequency boundary element method for scattering by a class of nonconvex obstacles*, Numer. Math., 129 (2015), pp. 647–689.
- [23] S. N. CHANDLER-WILDE AND S. LANGDON, *A Galerkin boundary element method for high frequency scattering by convex polygons*, SIAM J. Numer. Anal., 45 (2007), pp. 610–640.
- [24] F. COLLINO AND P. MONK, *The perfectly matched layer in curvilinear coordinates*, SIAM J. Sci. Comput., 19 (1998), pp. 2061–2090.
- [25] D. COLTON AND R. KRESS, *Inverse Acoustic and Electromagnetic Scattering Theory*, 2nd ed., Springer, New York, 1998.
- [26] D. L. COLTON AND R. KRESS, *Integral Equation Methods in Scattering Theory*, Wiley, New York, 1983.
- [27] J. B. CONWAY, *Functions of One Complex Variable I*, Springer, New York, 1978.
- [28] J.-P. CROISILLE AND G. LEBEAU, *Diffraction by an Immersed Elastic Wedge*, Lecture Notes in Math. 1723, Springer-Verlag, Berlin, 1999.
- [29] P.-M. CUTZACH AND C. HAZARD, *Existence, uniqueness and analyticity properties for electromagnetic scattering in a two-layered medium*, Math. Methods Appl. Sci., 21 (1998), pp. 433–461.
- [30] *NIST Digital Library of Mathematical Functions*, Release 1.0.28, F. W. J. Olver et al., eds., 2015, <http://dlmf.nist.gov/>.
- [31] S. DYATLOV AND M. ZWORSKI, *Mathematical Theory of Scattering Resonances*, Grad. Stud. Math. 200, AMS, Providence, RI, 2019.
- [32] S. FLISS AND P. JOLY, *Exact boundary conditions for time-harmonic wave propagation in locally perturbed periodic media*, Appl. Numer. Math., 59 (2009), pp. 2155–2178.
- [33] S. FLISS AND P. JOLY, *Solutions of the time-harmonic wave equation in periodic waveguides: Asymptotic behaviour and radiation condition*, Arch. Ration. Mech. Anal., 219 (2016), pp. 349–386.
- [34] S. FLISS, P. JOLY, AND J. LI, *Exact boundary conditions for wave propagation in periodic media containing a local perturbation*, in Wave Propagation in Periodic Media: Analysis, Numerical Techniques and Practical Applications, Progr. Comput. Phys. 1, Bentham, Sharjah, 2010, pp. 108–134.
- [35] D. P. HEWETT, S. L. LANGDON, AND J. M. MELENK, *A high frequency hp boundary element method for scattering by convex polygons*, SIAM J. Numer. Anal., 51 (2013), pp. 629–653.
- [36] C. JEREZ-HANCKES AND J.-C. NÉDÉLEC, *Asymptotics for Helmholtz and Maxwell solutions in 3-D open waveguides*, Commun. Comput. Phys., 11 (2012), pp. 629–646.

- [37] V. V. KAMOTSKIĬ, *An application of the method of spectral functions to the problem of scattering by two wedges*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI), 324 (2005), pp. 61–76 (in Russian); J. Math. Sci., 138 (2006), pp. 5514–5523, (in English).
- [38] N. KIELBASIEWICZ AND E. LUNÉVILLE, *User Documentation for XLiFE++*, 2019, https://uma.ensta-paris.fr/soft/XLiFE++/var/files/docs/usr/user_documentation.pdf.
- [39] W. LU, Y. Y. LU, AND J. QIAN, *Perfectly-matched-layer boundary integral equation method for wave scattering in a layered medium*, SIAM J. Appl. Math., 78 (2018), pp. 246–265.
- [40] W. MCLEAN, *Strongly Elliptic Systems and Boundary Integral Equations*, Cambridge University Press, Cambridge, UK, 2000.
- [41] I. MITREA, *On the spectra of elastostatic and hydrostatic layer potentials on curvilinear polygons*, J. Fourier Anal. Appl., 8 (2002), pp. 443–487.
- [42] J. OTT, *Halfspace Matching: A Domain Decomposition Method for Scattering by 2D Open Waveguides*, Ph.D. thesis, Karlsruhe Institute of Technology, 2017, <https://doi.org/10.5445/IR/1000070898>.
- [43] K.-M. PERFEKT AND M. PUTINAR, *The essential spectrum of the Neumann-Poincaré operator on a domain with corners*, Arch. Ration. Mech. Anal., 223 (2017), pp. 1019–1033.
- [44] S. A. SAUTER AND C. SCHWAB, *Boundary Element Methods*, Springer, New York, 2011.
- [45] Y. TJANDRAWIDJAJA, *Some Contributions to the Analysis of the Half-Space Matching Method for Scattering Problems and Extension to 3D Elastic Plates*, Ph.D. thesis, Université Paris Saclay, 2019.
- [46] A. TONNOIR, *Conditions transparentes pour la diffraction d'ondes en milieu élastique anisotrope*, Ph.D. thesis, École Polytechnique, 2015.